

Pancytopenia and isolated cytopenias as hematological complications of SLE: A narrative review

Oliwia Kubiś^{A–F}, Katarzyna Wojakowicz^{B–D,F}, Zuzanna Rzepkowska^{B–F},
Kacper Siemek^{B–D,F}, Alicja Dąbrowska^{A–F}, Magdalena Szmyrka^{A,C–F}

Faculty of Medicine, Wrocław Medical University, Wrocław, Poland

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation;
D – writing the article; E – critical revision of the article; F – final approval of the article

Advances in Clinical and Experimental Medicine, ISSN 1899–5276 (print), ISSN 2451–2680 (online)

Adv Clin Exp Med. 2026

Address for correspondence

Alicja Dąbrowska
E-mail: alicja.dabrowska@student.umw.edu.pl

Funding sources

None declared

Conflict of interest

None declared

Received on August 27, 2025
Reviewed on November 2, 2025
Accepted on November 21, 2025

Published online on September 14, 2026

Abstract

Pancytopenia is a significant hematological complication in patients with systemic lupus erythematosus (SLE), with an estimated prevalence of 10–40%. It is defined as a simultaneous decrease in red blood cells, white blood cells, and platelets. Although isolated cytopenias are more common, pancytopenia may indicate severe disease activity or secondary complications, such as bone marrow suppression or hemophagocytic syndromes. This review aims to summarize and synthesize current knowledge on the etiology, pathophysiology, diagnostic approach, and treatment strategies for pancytopenia in SLE, as well as its differentiation from other causes of bone marrow failure. The pathogenesis of pancytopenia in SLE is multifactorial and includes drug-induced bone marrow suppression, hypersplenism, myelofibrosis, macrophage activation syndrome (MAS), and autoimmune bone marrow failure. The diagnostic evaluation includes hematologic assessment, bone marrow examination, and exclusion of alternative diagnoses, such as aplastic anemia and paroxysmal nocturnal hemoglobinuria. Therapeutic management depends on the underlying cause. We also summarize published case reports comparing treatment approaches and clinical outcomes. The role of rituximab (RTX) in the management of pancytopenia and isolated cytopenias associated with SLE is discussed in detail. Particular attention is given to its mechanisms of action, safety profile and mixed clinical outcomes, documented in multicenter retrospective cohort studies, meta-analyses, and case series. Early recognition of pancytopenia remains a clinical challenge and requires well-structured diagnostic and therapeutic strategies. A thorough understanding of its underlying mechanisms and clinical manifestations is essential to avoid delays in treatment and prevent complications.

Key words: systemic lupus erythematosus, pancytopenia, rituximab, leukopenia, thrombocytopenia

Cite as

Kubiś O, Wojakowicz K, Rzepkowska Z, Siemek K, Dąbrowska A, Szmyrka M. Pancytopenia and isolated cytopenias as hematological complications of SLE: A narrative review [published online as ahead of print on September 14, 2026]. *Adv Clin Exp Med*. 2026. doi:10.17219/acem/214585

DOI

10.17219/acem/214585

Copyright

Copyright by Author(s)
This is an article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) (<https://creativecommons.org/licenses/by/3.0/>)

Highlights

- Pancytopenia affects 10–40% of patients with systemic lupus erythematosus (SLE) and may indicate severe disease activity or secondary bone marrow pathology.
- The pathogenesis is multifactorial, involving autoimmune bone marrow suppression, drug toxicity, hypersplenism, myelofibrosis, and macrophage activation syndrome.
- Accurate diagnosis requires comprehensive hematologic evaluation, bone marrow examination, and exclusion of differential diagnoses such as aplastic anemia and paroxysmal nocturnal hemoglobinuria.
- Rituximab demonstrates high initial response rates in SLE-related cytopenias, however randomized controlled trials show frequent relapses and difficulties maintaining long-term remissions.

Introduction

Systemic lupus erythematosus (SLE) is a chronic autoimmune disorder characterized by the production of autoantibodies against nuclear antigens and the formation of immune complexes that contribute to multiorgan tissue damage. Hematological manifestations are among the most common clinical features of SLE, with pancytopenia – defined as a concurrent reduction in red blood cells, white blood cells, and platelets – representing a particularly important clinical challenge. Its clinical presentation ranges from asymptomatic or mild, incidentally detected cases to severe manifestations associated with, e.g., sepsis.¹

Pancytopenia in the context of SLE may result from several pathogenic mechanisms, including autoimmune destruction of hematopoietic cells, hypersplenism, bone marrow suppression, and adverse effects of immunosuppressive therapy. Coexisting conditions, such as macrophage activation syndrome (MAS) or secondary myelofibrosis, may further complicate the clinical picture and pose significant diagnostic challenges.² Differentiating primary autoimmune cytopenias from other causes often requires a comprehensive diagnostic workup, including bone marrow biopsy, immunologic studies, and infectious disease screening.¹ The search for optimal diagnostic algorithms, supported by multicenter retrospective cohort studies, has led to the adoption of tools such as the Hemophagocytic Syndrome Diagnostic Score (HScore), which facilitates the diagnosis of hemophagocytic lymphohistiocytosis (HLH) and reduces the risk of overlooking the underlying cause of the patient's condition.³

Pancytopenia is considered not only a hematological manifestation of SLE but also a marker of disease activity and severity. Although isolated leukopenia, lymphopenia, and thrombocytopenia are relatively common, their coexistence as pancytopenia warrants a thorough etiological investigation. Effective treatment depends on identifying the underlying cause – whether uncontrolled autoimmunity, iatrogenic effects, or overlapping syndromes – and determining whether the pathology involves peripheral destruction of blood cells, intrinsic bone marrow dysfunction, or both.¹

Moreover, the 2nd part of this review focuses on the role of rituximab (RTX) in the treatment of pancytopenia and isolated cytopenias associated with SLE. Particular emphasis is placed on the mechanisms of action of RTX, with consideration given to the pathophysiological basis of the autoimmune process and its molecular targets. The efficacy of RTX is discussed in the context of evidence from multicenter retrospective cohort studies, meta-analyses, and case series. Its safety profile is also reviewed, with particular attention to patients with immune-mediated cytopenias.

Objectives

This review synthesizes current knowledge on the pathophysiology, differential diagnosis, and treatment of pancytopenia in SLE.⁴ Based on clinical studies, retrospective analyses, and current therapeutic guidelines, it aims to provide a comprehensive overview of this multifaceted complication while emphasizing the importance of an individualized, multidisciplinary approach.

Materials and methods

This review aimed to synthesize current knowledge on pancytopenia and other hematological complications associated with SLE. To achieve this, a comprehensive literature search was conducted using major scientific databases, focusing primarily on studies published within the past 15 years. The search keywords included “systemic lupus erythematosus,” “pancytopenia,” “cytopenia,” “autoimmune myelofibrosis,” “hemophagocytic syndrome,” “rituximab in SLE,” and “iatrogenic cytopenia.” We included studies addressing the pathophysiology, clinical presentation, differential diagnosis, and treatment of pancytopenia, as well as isolated cytopenias in the context of SLE. Eligible publications included original clinical studies, case series, and systematic reviews.

A total of more than 90 articles were screened, of which 71 peer-reviewed publications and case reports were

selected for in-depth analysis. These publications covered topics such as autoimmune mechanisms, drug-induced toxicity (e.g., iberdomide-induced degradation of the transcription factors Ikaros and Aiolos), laboratory biomarkers, differential diagnosis, and various immunosuppressive treatment protocols, including those described in case series as practical references for clinicians. In addition, the role of RTX in the treatment of immune-mediated cytopenias associated with SLE was thoroughly analyzed. Population-based studies were also included, providing broader insights into the true prevalence of hematological complications in this patient population. The data were organized thematically to illustrate the relationship between SLE and pancytopenia, leukopenia – with particular emphasis on neutropenia – and thrombocytopenia, followed by a section on differential diagnosis, including aplastic anemia, and concluding with evidence-based treatment strategies.

Pancytopenia in the context of SLE

Pancytopenia, defined as a reduction in all 3 major blood cell lineages, is a potential hematological complication in patients with SLE. This condition is diagnosed when hemoglobin (Hb) levels are below 10 g/dL, the total white blood cell count is $\leq 4 \times 10^9/L$, the neutrophil count is $< 1.5 \times 10^9/L$, and the platelet count is $< 100 \times 10^9/L$.⁵ Pancytopenia occurs less frequently than isolated cytopenias in patients with SLE. It is estimated to affect 10–40% of patients with SLE, whereas neutropenia and thrombocytopenia are observed more frequently than aplastic anemia.^{5,6}

Pathogenesis of pancytopenia in SLE

Potential causes of pancytopenia in patients with SLE include medications, peripheral blood cell destruction, autoimmune myelofibrosis, and rare conditions such as MAS.⁷ In addition, bone marrow aspiration typically reveals hypocellularity and necrosis, which are attributed to autoimmune mechanisms. Various triggering factors, including immune activation and infections, may lead to excessive macrophage activation in the bone marrow and other tissues.⁸ Moreover, hemophagocytosis is frequently observed in the bone marrow but is neither pathognomonic nor sufficient for establishing the diagnosis.⁹

Autoimmune diseases such as SLE, rheumatoid arthritis, and Felty's syndrome contribute to pancytopenia through hypersplenism, which results in peripheral destruction of blood cells by the reticuloendothelial system. Furthermore, hypersplenism occurs in 9–46% of patients with SLE,¹⁰ particularly during periods of active disease.⁵

Myelofibrosis is another important cause of pancytopenia and is characterized by clonal proliferation of myeloid stem cells and the production of a fibrotic bone marrow

stroma.⁶ In addition, life-threatening conditions such as HLH, disseminated intravascular coagulation (DIC), and paroxysmal nocturnal hemoglobinuria (PNH) should also be considered in the differential diagnosis of pancytopenia in critically ill patients.⁵

Iatrogenic pancytopenia in SLE

Pancytopenia may be induced by various iatrogenic factors, including chemotherapy, radiotherapy, and antiepileptic drugs such as carbamazepine and valproate. Drug interactions, including those between allopurinol and azathioprine (AZA) or methotrexate (MTX) and trimethoprim, may also contribute to the development of pancytopenia. Additional causes include sepsis and certain antibiotics, particularly chloramphenicol, trimethoprim, and linezolid.⁵

In addition, immunosuppressive drugs used in the treatment of SLE, including AZA, MTX, and cyclophosphamide (CYC), may cause bone marrow suppression.⁵ The mechanisms underlying drug-induced cytopenia include immune-mediated reactions. These reactions may impair hematopoiesis within the bone marrow or enhance peripheral destruction of blood cells. They may also be associated with the patient's immune profile or coexisting diseases.⁶ Moreover, pancytopenia in SLE may be induced by interferon alpha (IFN- α) in the context of HLH.¹¹ Cases of pancytopenia developing during sulfasalazine therapy have also been reported. Furthermore, resolution of pancytopenia following discontinuation of sulfasalazine has been described.¹²

Anemia

Studies consistently identify anemia as the most common hematological manifestation in patients with SLE, with a reported prevalence exceeding 50%.⁶ Another review reported rates as high as 98%, reflecting the diverse etiologies of anemia in SLE, including anemia of chronic disease, autoimmune hemolytic anemia, and microangiopathic hemolytic anemia. Anemia of chronic disease was the predominant subtype, underscoring the impact of chronic inflammation and hematologic dysregulation associated with SLE.¹³

Anemia of chronic disease is considered the most common type of anemia in SLE, accounting for approx. 33% of cases. In asymptomatic patients, treatment beyond adequate control of the underlying disease is usually unnecessary; however, in severe cases, erythropoiesis-stimulating agents may be considered.⁶

Autoimmune hemolytic anemia (AIHA) occurs in approx. 10% of patients with SLE, may precede the diagnosis of SLE by several years, and may represent the predominant clinical manifestation. Warm AIHA is the most common subtype, and its treatment is based primarily

on glucocorticosteroids (GCs). However, increasing attention has been paid to the use of RTX in patients with SLE-associated AIHA, as it may avoid the need for splenectomy and its associated complications.⁶ Aplastic anemia may also occur in patients with SLE, and careful monitoring while awaiting the response to treatment is essential because blood transfusions may be required.⁶

Microangiopathic hemolytic anemia leading to thrombotic microangiopathy (TMA) occurs in approx. 0.5–10% of patients with SLE and is associated with a poor clinical prognosis. Although this is a rare complication, its coexistence with SLE is associated with high mortality, ranging from 33.9% to 62.5%, despite the use of plasmapheresis, which remains the cornerstone of treatment in this setting.⁶

Regardless of the underlying etiology of anemia in SLE, it is essential to determine whether the patient requires urgent intervention, including blood transfusion. It should be emphasized that transfusion is a supportive measure aimed at stabilizing the patient's condition until immunosuppressive or hematological treatment becomes effective.

Transfusion is generally considered when the Hb concentration is <7 g/dL in hemodynamically stable patients, or <9 g/dL in patients with cardiovascular disease (CVD), impaired tissue perfusion, acute hemolytic anemia, or active AIHA. Transfusion is also indicated in symptomatic AIHA accompanied by signs of circulatory failure and in anemia resulting from aplastic anemia or myelofibrosis when causal treatment is ineffective.¹

In patients with SLE, the use of leukoreduced blood components is recommended because it reduces the risk of febrile nonhemolytic transfusion reactions, alloimmunization, and transmission of leukocyte-associated viruses. In addition, irradiated blood components are recommended for patients with profound immunosuppression, particularly those receiving RTX, CYC, mycophenolate mofetil, or those who have undergone hematopoietic stem cell transplantation (HSCT). This approach aims to prevent transfusion-associated graft-versus-host disease (TA-GVHD), a potentially life-threatening complication in this patient population.¹

Leukopenia

Leukopenia, defined as a reduced white blood cell count, is a common hematological manifestation of SLE and often correlates with disease activity. Proposed mechanisms include increased peripheral destruction of granulocytes, alterations in the marginal granulocyte pool and spleen, and reduced bone marrow production.⁶ The most common form of lymphopenia in SLE is T-cell lymphopenia, which is strongly associated with disease activity, particularly when accompanied by thrombocytopenia. Notably, both the American College of Rheumatology (ACR) and

the Systemic Lupus International Collaborating Clinics (SLICC) recognize leukopenia as an important classification criterion for SLE.⁶

In addition, lymphopenia has been reported to be significantly associated with overall SLE disease activity, erythrocyte sedimentation rate (ESR), serological findings, and the use of various medications, including prednisolone, AZA, MTX, CYC, and RTX.¹⁴ In contrast, MTX and cyclosporine have been reported to be associated with a lower risk of neutropenia. Lower SLE disease activity has likewise been associated with a lower prevalence of both lymphopenia and neutropenia.¹⁵

Neutropenia

Neutropenia is the most common neutrophil abnormality in SLE, occurring in 20–47% of patients. It is classified as mild when the neutrophil count is $1.0\text{--}1.5 \times 10^9/\text{L}$, moderate at $0.5\text{--}1.0 \times 10^9/\text{L}$, and severe at $<0.5 \times 10^9/\text{L}$.¹⁶ Neutrophils have long been recognized as a potential source of autoantigens and inflammatory cytokines in SLE.¹⁷ Neutrophil populations in patients with SLE differ from those in healthy individuals, with impaired phagocytic function, altered metabolism, and increased apoptosis representing the principal abnormalities.¹⁸ Consequently, these neutrophils exhibit an increased capacity to form neutrophil extracellular traps (NETs) and contain autoantigens as well as elevated levels of other immunostimulatory molecules.¹⁹ Excessive NET formation may contribute to disease progression by stimulating type I interferon production by plasmacytoid dendritic cells and promoting endothelial dysfunction and prothrombotic changes.¹⁷

In addition to reduced bone marrow production and immune-mediated destruction, transient reductions in circulating neutrophil counts in SLE may also result from increased margination of neutrophils along the vascular endothelium.²⁰ Margination reflects the redistribution of neutrophils from the freely circulating pool into marginated intravascular reservoirs (e.g., the spleen and the pulmonary and hepatic microvasculature) and may result in apparent peripheral neutropenia without a true reduction in neutrophil mass.²¹ Moreover, iberdomide, a novel therapeutic agent under investigation for SLE, promotes degradation of the transcription factors Ikaros and Aiolos, thereby influencing leukocyte development and autoimmune processes. However, its use is associated with adverse effects, including neutropenia.²²

A study comparing 208 patients with neutropenia and 779 patients with SLE without neutropenia found that neutropenia was significantly associated with thrombocytopenia, lymphopenia, and reduced complement C3 levels.¹⁷ Chronic neutropenia was also associated with the presence of anti-Ro/SSA antibodies. Kurien et al. proposed that anti-Ro/SSA antibodies may cross-react with neutrophil surface

proteins, triggering complement activation and subsequent cell damage.²³ Furthermore, Beyan et al.²⁴ reported a 20% prevalence of neutropenia among 115 patients with SLE ($<1.8 \times 10^9/L$), whereas Dias et al.²⁵ documented a prevalence of 40.3% among 124 patients with SLE, with severe and persistent neutropenia occurring in 0.8% and 4.8% of patients respectively.

Thrombocytopenia

Mild thrombocytopenia (platelet count: $100\text{--}150 \times 10^9/L$) is observed in 25–50% of patients, whereas severe thrombocytopenia (platelet count $<50 \times 10^9/L$) occurs in approx. 10%.²⁶ Immune thrombocytopenia (ITP), the primary cause of SLE-associated thrombocytopenia,⁶ is defined as a platelet count $<100 \times 10^9/L$ and affects 10–40% of patients with SLE.²⁷ Immune thrombocytopenia may precede the clinical manifestations of SLE in 5–16% of patients.²⁸ Moreover, severe thrombocytopenia is associated with a poor prognosis and increased mortality.²⁹ The underlying mechanisms include impaired platelet production by megakaryocytes and increased peripheral platelet destruction mediated by autoimmune processes.⁷

Treatment of SLE-associated immune thrombocytopenia

Currently, the treatment of SLE-associated ITP (SLE-ITP) is based on immunosuppressive therapy, similarly to the treatment of primary ITP. Approximately 75% of patients with SLE-ITP respond to GCs and antimalarial drugs.²⁹ In cases in which ITP occurs secondary to SLE, GCs play a central role in treatment, resulting in clinical improvement in approx. 2/3 of patients. Moreover, GC monotherapy achieves remission in nearly 20% of patients. Dexamethasone is the preferred GC because of its more rapid onset of action and lower risk of toxicity compared with prolonged prednisolone therapy.³⁰ Furthermore, intravenous immunoglobulin (IVIG) can temporarily increase platelet counts; however, it is used primarily in patients who are intolerant of corticosteroids. Immunoglobulin G (IgG) is administered intravenously at a dose of 1 g/kg/day for 1 or 2 days.⁶

The latest therapeutic options include thrombopoietin receptor agonists (TPO-RAs), platelet desialylation inhibitors (e.g., oseltamivir),³¹ neonatal Fc receptor (FcRn) inhibitors, spleen tyrosine kinase (Syk) inhibitors, Bruton's tyrosine kinase (BTK) inhibitors,³² and B-cell-targeted therapies (e.g., RTX and belimumab).³³

The introduction of RTX into the treatment of ITP has resulted in response rates of up to 60%, providing an alternative to splenectomy. However, observational studies indicate that only 20–30% of patients maintain long-term remission, with treatment being more effective in younger women before the disease enters the chronic phase.³⁴

Differentiation of pancytopenia in SLE from aplastic anemia and other types of anemia

The diagnosis of aplastic anemia (AA), a rare complication in patients with SLE, is often delayed because of the presence of other potential causes of pancytopenia.³⁵ In addition, several other types of anemia may occur in patients with SLE, including microangiopathic hemolytic anemia, iron deficiency anemia, autoimmune hemolytic anemia (with a positive Coombs test), pure red cell aplasia, and anemia secondary to chronic kidney disease (CKD). Autoimmune cytopenias in SLE are associated with the presence of autoantibodies directed against blood cell antigens, leading to immune-mediated destruction of hematopoietic cells.⁴

Differential diagnosis of pancytopenia in SLE

Once pancytopenia has been confirmed based on the initial complete blood count, preliminary noninvasive investigations are recommended, including reticulocyte count and peripheral blood smear analysis. The reticulocyte count serves as a reliable indicator of bone marrow function.¹⁴

In recent years, several biomarkers of primary myelofibrosis have been identified, including circulating YKL-40 and GATA-1; however, they are not useful for diagnosing myelofibrosis associated with SLE.³⁶ Further hematological evaluation is typically recommended, including bone marrow aspiration and trephine biopsy, which may reveal a hypercellular bone marrow. Trephine biopsy is also used to confirm hypocellularity or bone marrow necrosis⁵ (Fig. 1).

Chronic SLE and long-term immunosuppressive therapy increase the risk of lymphoid malignancies. Therefore, in patients with persistent cytopenia or new clinical features, such as progressive lymphadenopathy, fever, night sweats, or weight loss, lymphoma should be considered in the differential diagnosis.⁵

In the differential diagnosis, it is important to distinguish between bone marrow suppression caused by medications or active autoimmune disease and malignant bone marrow infiltration. The final diagnosis is based on bone marrow aspiration and trephine biopsy with histopathological evaluation, flow cytometry, and cytogenetic and molecular analyses for monoclonal lymphocyte populations, allowing lymphoma to be confirmed or excluded as the cause of pancytopenia. In diagnostically challenging cases, additional molecular and immunohistochemical analyses, including GATA-1 assessment, may be considered to help distinguish myeloproliferative neoplasms from lymphoproliferative disorders.⁵

Moreover, infectious causes should also be considered in the differential diagnosis of pancytopenia in patients




Fig. 1. Pathogenesis and differential diagnosis of pancytopenia in systemic lupus erythematosus (SLE)

RA – rheumatoid arthritis; MAS – macrophage activation syndrome; DIC – disseminated intravascular coagulation; PNH – paroxysmal nocturnal hemoglobinuria; HLH – hemophagocytic lymphohistiocytosis; AST – aspartate aminotransferase; $TNF-\alpha$ – tumor necrosis factor alpha.

with SLE. Viral infections, including parvovirus B19, HIV, Epstein–Barr virus (EBV), cytomegalovirus (CMV), and hepatitis viruses (HAV, HBV, and HCV), may impair bone marrow hematopoiesis because of their tropism for hematopoietic progenitor cells and their ability to induce aplastic anemia.⁵ In endemic regions, visceral leishmaniasis should also be considered because it may closely mimic SLE flares by presenting with fever, pancytopenia, splenomegaly, hypergammaglobulinemia, and autoantibody production.³⁷

Furthermore, additional investigations, including flow cytometry for PNH and measurements of aspartate aminotransferase (AST), ferritin, triglycerides, and fibrinogen, are important for calculating the HScore in patients with suspected HLH.⁵

Treatment of pancytopenia is primarily directed at identifying and eliminating the underlying cause (Table 1). In addition, antibiotics are indicated in cases of severe neutropenia, and blood transfusions may be administered when clinically indicated to manage severe anemia.⁵ Prompt recognition and comprehensive diagnostic evaluation are essential to distinguish SLE-related pancytopenia from other potentially life-threatening conditions.

A thorough understanding of the pathophysiology and clinical implications of pancytopenia in SLE is crucial for guiding appropriate therapeutic strategies and improving patient outcomes. Further research is warranted to clarify the underlying mechanisms and optimize management strategies for this complex and heterogeneous patient population.

The role of rituximab in the treatment of cytopenias associated with SLE

Given the central role of B lymphocytes in the pathogenesis of SLE, RTX, a chimeric mouse/human monoclonal antibody targeting the CD20 antigen, has been evaluated in numerous randomized controlled trials for the treatment of SLE, including studies assessing its B-cell specificity and efficacy in SLE as well as in other rheumatic diseases.⁴⁸ Rituximab induces depletion of mature B lymphocytes through complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC),

Table 1. Summary of case reports on systemic lupus erythematosus (SLE)-associated pancytopenia and isolated cytopenias

Authors and year of publication	Case report	Treatment	Clinical outcome
Serin et al., 2021 ³⁸	30-year-old female patient with SLE, suspected aplastic anemia/myelodysplastic syndrome (MDS) with a positive fluorescein-labeled aerolysin (FLAER) test.	Pulse methylprednisolone 1 g/day for 2 days; prednisone 20 mg; cyclosporine 2 × 50 mg.	Good response; transfusion no longer needed; patient transferred to outpatient care.
Frikha et al., 2022 ³⁹	45-year-old woman with rheumatoid arthritis, agranulocytosis in SLE.	Methylprednisolone pulses (1 g/day × 3 days); hydroxychloroquine 400 mg/day; adjuvant iron and folic acid; high-dose corticosteroids (1 mg/kg/day) × 8 weeks.	Resolution of epigastralgia, fever, diarrhea, leukopenia, and thrombocytopenia.
Sahbani et al., 2024 ⁴⁰	16-year-old boy with hemorrhagic syndrome (atypical presentation of SLE).	Dexamethasone 40 mg/day × 3 days; intravenous immunoglobulin (IVIG) 0.5 g/kg × 4 days; platelet transfusion; later hydroxychloroquine and corticosteroids; maintenance with azathioprine (AZA) and corticosteroids.	Stabilization of thrombocytopenia; long-term stability on AZA and corticosteroids.
Horino et al., 2025 ⁴¹	67-year-old woman with pancytopenia and schizophrenia (neuropsychiatric SLE in a patient with pancytopenia and chronic schizophrenia).	Oral prednisolone 40 mg/day, tapered to 5 mg/day over 2 years.	Improvement in pancytopenia, pleural effusion, and ascites; no recurrence during 2-year follow-up.
Haidar et al., 2024 ⁴²	63-year-old female patient with pancytopenia as a first presentation of late-onset SLE.	Prednisone 60 mg/day; hydroxychloroquine 200 mg/day; prednisolone taper 5 mg/week.	Improved fatigue, no fever; normalization of white blood cells, hemoglobin, and platelets; decreased erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP); 2 kg weight gain in 4 weeks.
Brar et al., 2022 ⁴³	73-year-old male patient, hydralazine-induced lupus presenting with pancytopenia.	Discontinuation of hydralazine.	Marked symptom improvement; complete resolution of night sweats, fevers, and weight loss within 2 months.
Ungprasert et al., 2016 ⁴⁴	33-year-old female patient with autoimmune myelofibrosis with pancytopenia as a presenting manifestation of SLE.	Glucocorticoids; mycophenolate mofetil.	Good response to therapy.
Guevara et al., 2023 ⁴⁵	34-year-old female patient with pancytopenia and autoimmune history.	Steroids; hydroxychloroquine.	Complete resolution of symptoms; pancytopenia returned to baseline with mild persistent anemia.
Abu Jheasha et al., 2024 ⁴⁶	22-year-old female patient with protein-losing enteropathy (rare SLE manifestation).	Pulse methylprednisolone 1 g/day; hydroxychloroquine 200 mg/day.	Clinical and laboratory improvement; stable vitals; discharged with regular rheumatology follow-up.
Ameer et al., 2024 ⁴⁷	30-year-old female patient with pancytopenia and SLE.	Mycophenolate 1,500 mg twice daily; lisinopril 30 mg/day; prednisone 10 mg/day; ongoing acute promyelocytic leukemia (APML) therapy with all-trans-retinoic acid/arsenic trioxide (ATRA/ATO).	Continuation of therapy until remission.

and induction of apoptosis, providing a biological rationale for its use in autoimmune cytopenias associated with SLE.⁴⁹

In many countries, RTX is used off-label for the treatment of primary immune cytopenias in adults. Prospective randomized studies have demonstrated response rates of approx. 50–60% in ITP⁵⁰ and up to 75% in warm AIHA.⁵¹ However, studies evaluating RTX in SLE-related cytopenias have not demonstrated a significant clinical benefit; therefore, its use in this setting remains off-label.⁴⁸

Mechanisms of action of RTX

Direct depletion of CD20-positive B lymphocytes through CDC and ADCC prevents the generation of autoantibody-producing plasma cells directed against blood

cell antigens (e.g., antiplatelet antibodies in ITP and antierythrocyte antibodies in AIHA), thereby reducing opsonization and peripheral destruction of blood cells.⁴ In cases of severe thrombocytopenia, RTX treatment stabilized platelet counts above $100 \times 10^9/\mu\text{L}$ for more than 6 months and was accompanied by the disappearance of anti-DNA antibodies⁴ (Fig. 2).

Importantly, RTX targets both immature pre-B cells in the bone marrow and mature circulating B lymphocytes. As B cells differentiate into immunoglobulin-producing plasma cells, they lose CD20 expression. Consequently, plasma cells are generally not depleted during RTX therapy, and serum immunoglobulin levels usually remain within the normal range. In addition, RTX does not affect hematopoietic stem cells (HSCs).⁵²




Fig. 2. Mechanism of action of rituximab (RTX)

CDC – complement-dependent cytotoxicity; ADCC – antibody-dependent cellular cytotoxicity; FcγR – Fc gamma receptor; C1q – complement component 1q; RBC – red blood cell; PLT – platelet; Tfh – T follicular helper cell; Tfr – T follicular regulatory cell.

B-cell depletion also influences the number and function of helper T lymphocytes, including T follicular helper (Tfh) cells, as well as cytokine production, which is relevant in immune-mediated bone marrow suppression.³³ Furthermore, RTX exerts complement-dependent cytotoxicity by binding C1q, leading to the formation of the membrane attack complex and subsequent lysis of CD20-positive B cells. In addition, RTX induces apoptosis through activation of the caspase-3 signaling pathway and via Fab-dependent mechanisms.⁵³ Antibody-dependent cellular cytotoxicity represents another important mechanism of RTX action. This process involves binding of Fc receptors expressed on monocytes, macrophages, and natural killer (NK) cells to the Fc region of RTX, resulting in activation of effector cells and lysis of B lymphocytes.⁵⁴ Rituximab may also modulate germinal center responses and the activity of follicular regulatory T (Tfr) cells, which suppress B-cell activation within germinal centers.⁵³

In cases associated with MAS or HLH, the therapeutic effect of RTX may result from disruption of the pathological feedback loop between autoreactive B cells and macrophage

overactivation through modulation of the cytokine network. Rituximab may be used both as monotherapy and as an adjunct to the HLH-2004 treatment protocol. Compared with etoposide, RTX has a more favorable safety profile and is generally better tolerated.⁵⁵

Efficacy of RTX in the treatment of hematological manifestations of SLE

Evidence supporting the efficacy of RTX in the treatment of hematological manifestations of SLE is derived primarily from case series, large observational cohorts, and meta-analyses. In a multicenter retrospective study by Serris et al., RTX produced a high rate of rapid responses in immune cytopenias associated with SLE. The overall initial response rate was 82%, including a complete response (CR) rate of 59%. The initial response rates were 87% for ITP, 86% for AIHA, and 50% for Evans syndrome. Although relapse occurred in 41% of patients despite an initial response, re-treatment with RTX was effective in 94% of these cases.⁵⁶

In the study by Sans-Pola et al., RTX was administered to 33 patients with extrarenal SLE, resulting in partial or complete responses. The median SLEDAI-2K score decreased from 9 to 1.5, and patients with thrombocytopenia experienced significant increases in platelet counts. Anti-double-stranded DNA (anti-dsDNA) antibody levels also declined, from a median of 64.3 to 32.7. However, the median time to relapse after the first treatment cycle was 1.6 years, and all patients required ongoing therapy to maintain remission.⁵⁷

In a multicenter retrospective cohort study including 71 adults with SLE-associated cytopenias, 86% of patients responded initially to RTX (91% with ITP, 87.5% with AIHA, and 60% with Evans syndrome), and 60.5% achieved complete remission. No cases of RTX-induced neutropenia were reported, and only 3 patients developed severe infections. Among the 24 patients (39.3%) who relapsed after an initial response, 18 underwent re-treatment with RTX, which was successful in 88.8% of cases.⁴⁹

Pediatric case reports have also described severe, treatment-refractory pancytopenia that achieved remission following RTX therapy without significant adverse events.⁵⁸

Although the placebo-controlled randomized EXPLORER (extrarenal lupus) and LUNAR (lupus nephritis) trials did not demonstrate statistically significant superiority of RTX combined with standard therapy, observational evidence suggests that carefully selected patients with hematological manifestations of SLE, including ITP, AIHA, and pancytopenia, may benefit from RTX treatment.⁵⁹

In an interventional study by Zhang et al., 8 patients with treatment-refractory SLE-associated ITP received low-dose RTX (2 infusions of 200 mg administered at 2-week intervals) together with tapering corticosteroid therapy. Overall response rates (ORRs) at 1, 3, 6, and 12 months were 25.0%, 87.5%, 75.0%, and 75.0%, respectively, suggesting that this regimen may represent a promising therapeutic approach.⁶⁰

In a retrospective cohort study by Harris et al., the ORR to RTX (56%) did not differ significantly between children (58%, $n = 24$) and adults (55%, $n = 40$). Response rates were also similar in primary and secondary ITP (53% and 62%, respectively). Evans syndrome was more common in children (42%) than in adults (18%) treated with RTX.⁶¹

Safety profile of RTX, with emphasis on patients with immune cytopenias

Rituximab is not without adverse effects. It may induce secondary hypogammaglobulinemia, thereby increasing the risk of severe infections, particularly in patients with low IgG levels before treatment or after repeated treatment courses. This complication may require IVIG replacement therapy. The risk is particularly relevant in patients with baseline pancytopenia, as this condition itself predisposes them to infections and necessitates regular monitoring of serum immunoglobulin levels.⁶²

In addition, late-onset neutropenia (LON) may occur weeks to months after RTX administration, although it is usually transient. Concomitant use of other immunosuppressive agents or previous treatment courses may increase this risk.⁶³

Infusion-related reactions may also occur; however, they are generally mild to moderate.⁵³ There are also isolated reports suggesting that RTX may exacerbate bone marrow suppression.⁶⁴

Other treatment methods in SLE patients with cytopenias

Clinical studies have demonstrated that mesenchymal stem cell (MSC) transplantation can improve outcomes in patients with SLE by enhancing regulatory T-cell (Treg) and T-helper 2 (Th2) responses while suppressing pro-inflammatory Th1, Th17, and B-cell responses, with an overall favorable safety profile. However, therapeutic efficacy remains variable, which may reflect differences in donor cell quality and recipient immune characteristics.⁶⁵

An alternative treatment for severe SLE is hematopoietic stem cell transplantation (HSCT), which may induce long-term remission but is associated with risks such as infections and conditioning-related toxicity.⁶⁶

Belimumab, a B-lymphocyte stimulator (BLyS/BAFF) inhibitor, has beneficial effects on the course of SLE, including its hematological manifestations, by suppressing excessive activation of autoreactive B lymphocytes.²⁹ Analyses of published case reports suggest that belimumab may improve platelet counts and Hb concentrations in patients with SLE-ITP and autoimmune hemolytic anemia (AIHA).⁶⁷ In addition, belimumab therapy may allow reduction of GC doses.⁶⁸ Its safety profile has been shown to be favorable across different clinical phenotypes of SLE. However, it should be noted that patients with severe, life-threatening cytopenias were excluded from the pivotal registration trials. Therefore, belimumab may be considered an adjunctive treatment option for chronic, refractory cytopenias associated with SLE, although the decision to initiate therapy should be individualized.⁶⁹

Another biologic agent that may be useful in the treatment of cytopenias in patients with SLE is anifrolumab. It is a monoclonal antibody that blocks the type I interferon receptor (IFNAR1), thereby targeting a key pathogenic pathway in SLE and reducing disease activity, including improvement of hematological parameters in some patients.⁷⁰ Post hoc analyses of clinical trials have shown that inhibition of type I interferon signaling is associated with improvements across multiple domains of disease activity, including hematological manifestations. However, no clinical studies have specifically evaluated anifrolumab for the treatment of severe cytopenias in patients with SLE.⁷¹

Limitations of the study

This review has several limitations, primarily related to the heterogeneity and limited availability of large, high-quality studies investigating pancytopenia in SLE. Much of the available evidence is derived from small case series, introducing the potential for selection and reporting bias. In addition, the retrospective design of many studies and the overlap between iatrogenic causes and disease manifestations limit causal inference. There is a clear need for prospective cohort studies and randomized clinical trials to support the development of robust diagnostic and therapeutic guidelines for the management of hematological disorders in patients with SLE.

Conclusions

Pancytopenia in SLE is a multifactorial complication with a diverse etiology. It may reflect primary autoimmune activity or arise secondary to infections, drug toxicity, bone marrow suppression, or concomitant hematological disorders such as MAS. Early recognition and a systematic diagnostic evaluation are essential for identifying the underlying cause and selecting appropriate treatment. Bone marrow examination, serological testing, and immunologic profiling are key components of the diagnostic workup and help exclude conditions that may mimic SLE-associated pancytopenia, such as hematological malignancies or aplastic anemia.

Therapeutic interventions should be directed at the underlying cause. In autoimmune-mediated pancytopenia, immunosuppressive therapy – including GCs, IVIG, and biologic agents such as RTX – may be effective. In cases related to drug toxicity or infection, supportive care and modification of the immunosuppressive regimen are required. Novel therapeutic approaches, including neonatal Fc receptor (FcRn) inhibitors and immunomodulatory agents such as iberdomide, are currently under clinical investigation and may offer promising treatment options for refractory cases.

Importantly, pancytopenia should not be regarded merely as a laboratory abnormality; it often reflects increased disease activity and an elevated risk of systemic complications. Its presence should prompt comprehensive clinical reassessment and multidisciplinary collaboration involving rheumatologists, hematologists, and infectious disease specialists.

Further research is needed to clarify the immunopathogenesis of pancytopenia in SLE and to optimize evidence-based treatment algorithms. Given its potential for rapid progression and its prognostic significance, pancytopenia should be recognized as a serious clinical manifestation requiring prompt evaluation and appropriate management within the broader context of systemic autoimmunity.

Use of AI and AI-assisted technologies

Not applicable.

ORCID iDs

Oliwia Kubiś  <https://orcid.org/0009-0005-2576-8432>
 Katarzyna Wojakowicz  <https://orcid.org/0009-0007-8812-9982>
 Zuzanna Rzepkowska  <https://orcid.org/0009-0007-1950-7302>
 Kacper Siemek  <https://orcid.org/0009-0001-8983-1411>
 Alicja Dąbrowska  <https://orcid.org/0000-0001-8610-107X>
 Magdalena Szmyrka  <https://orcid.org/0000-0001-6885-6153>

References

- Gnanaraj J, Parnes A, Francis CW, Go RS, Takemoto CM, Hashmi SK. Approach to pancytopenia: Diagnostic algorithm for clinical hematologists. *Blood Rev.* 2018;32(5):361–367. doi:10.1016/j.blre.2018.03.001
- Gouda W, Alsaqabi F, Moshirif A, Abbas AS, Abdel-Aziz TM, Islam MA. Macrophage activation syndrome triggered by systemic lupus erythematosus flare: Successful treatment with a combination of dexamethasone sodium phosphate, intravenous immunoglobulin, and cyclosporine. A case report. *J Med Case Rep.* 2021;15(1):497. doi:10.1186/s13256-021-03072-1
- Fardet L, Galicier L, Lambotte O, et al. Development and validation of the HScore, a score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis Rheumatol.* 2014;66(9):2613–2620. doi:10.1002/art.38690
- Bashal F. Hematological disorders in patients with systemic lupus erythematosus. *Open Rheumatol J.* 2013;7(1):87–95. doi:10.2174/1874312901307010087
- Chew S, Kamangar M. Approach to pancytopenia: From blood tests to the bedside. *Clin Med (Lond).* 2024;24(5):100235. doi:10.1016/j.clinme.2024.100235
- Santacruz JC, Mantilla MJ, Rueda I, Pulido S, Rodriguez-Salas G, Londono J. A practical perspective of the hematologic manifestations of systemic lupus erythematosus. *Cureus.* 2022;14(3):e22938. doi:10.7759/cureus.22938
- Velo-García A, Castro SG, Isenberg DA. The diagnosis and management of the haematologic manifestations of lupus. *J Autoimmun.* 2016;74:139–160. doi:10.1016/j.jaut.2016.07.001
- Wanitpongpan C, Teawtrakul N, Mahakkanukrauh A, Siritunyaporn S, Sirijerachai C, Chansung K. Bone marrow abnormalities in systemic lupus erythematosus with peripheral cytopenia. *Clin Exp Rheumatol.* 2012;30(6):825–829. PMID:23101438.
- Lambotte O, Khellaf M, Harmouche H, et al. Characteristics and long-term outcome of 15 episodes of systemic lupus erythematosus-associated hemophagocytic syndrome. *Medicine (Baltimore).* 2006;85(3):169–182. doi:10.1097/01.md.0000224708.62510.d1
- Cruz AJ, Castro A. Systemic lupus erythematosus presenting as spontaneous splenic rupture. *BMJ Case Rep.* 2015;2015:bcr2015121531. doi:10.1136/bcr-2015-212531
- Zeng Z, Tu W, Ji B, et al. IFN- α induced systemic lupus erythematosus complicated with hemophagocytic lymphohistiocytosis: A case report and literature review. *Front Immunol.* 2023;14:1223062. doi:10.3389/fimmu.2023.1223062
- Alanazi AS. Sulfasalazine-induced pancytopenia indicative of bone marrow suppression: A case report. *J Med Cases.* 2014;5(5):289–291. doi:10.14740/jmc1767w
- Bathina A, Chintada DC, Yellu NKR, Vijayashree J, Khatija Begum M, Unnikrishnan P. Clinical and hematological manifestations of systemic lupus erythematosus at initial presentation in a tertiary healthcare center. *Cureus.* 2024;16(12):e75956. doi:10.7759/cureus.75956
- Priya P, Subhashree AR. Role of absolute reticulocyte count in evaluation of pancytopenia: A hospital based study. *J Clin Diagn Res.* 2014;8(8):FC01–03. doi:10.7860/JCDR/2014/8949.4704
- Kandane-Rathnayake R, Louthrenoo W, Golder V, et al. Independent associations of lymphopenia and neutropenia in patients with systemic lupus erythematosus: A longitudinal, multinational study. *Rheumatology (Oxford).* 2021;60(11):5185–5193. doi:10.1093/rheumatology/keab217

16. Oliveira GHM, Nesbitt GC, Murphy JG, eds. *Mayo Clinic Medical Manual*. Boca Raton, USA: Mayo Clinic Scientific Press, Taylor & Francis Group; 2006. ISBN:978-0-8493-9087-6.
17. Meyer A, Guffroy A, Blaison G, et al. Systemic lupus erythematosus and neutropaenia: A hallmark of haematological manifestations. *Lupus Sci Med*. 2020;7(1):e000399. doi:10.1136/lupus-2020-000399
18. Wang H, Kim SJ, Lei Y, et al. Neutrophil extracellular traps in homeostasis and disease. *Sig Transduct Target Ther*. 2024;9(1):235. doi:10.1038/s41392-024-01933-x
19. Gestermann N, Di Domizio J, Lande R, et al. Netting neutrophils activate autoreactive B cells in lupus. *J Immunol*. 2018;200(10):3364–3371. doi:10.4049/jimmunol.1700778
20. Wigerblad G, Kaplan MJ. Neutrophil extracellular traps in systemic autoimmune and autoinflammatory diseases. *Nat Rev Immunol*. 2023;23(5):274–288. doi:10.1038/s41577-022-00787-0
21. De Filippo K, Rankin SM. The secretive life of neutrophils revealed by intravital microscopy. *Front Cell Dev Biol*. 2020;8:603230. doi:10.3389/fcell.2020.603230
22. Merrill JT, Werth VP, Furie R, et al. Phase 2 trial of iberdomide in systemic lupus erythematosus. *N Engl J Med*. 2022;386(11):1034–1045. doi:10.1056/NEJMoa2106535
23. Kurien BT, Newland J, Paczkowski C, Moore KL, Scofield RH. Association of neutropenia in systemic lupus erythematosus (SLE) with anti-Ro and binding of an immunologically cross-reactive neutrophil membrane antigen. *Clin Exp Immunol*. 2001;120(1):209–217. doi:10.1046/j.1365-2249.2000.01195.x
24. Beyan E, Beyan C, Turan M. Hematological presentation in systemic lupus erythematosus and its relationship with disease activity. *Hematology*. 2007;12(3):257–261. doi:10.1080/10245330701214145
25. Dias AMB, Do Couto MCM, Duarte CCM, Inês LPB, Malcata AB. White blood cell count abnormalities and infections in one-year follow-up of 124 patients with SLE. *Ann N Y Acad Sci*. 2009;1173(1):103–107. doi:10.1111/j.1749-6632.2009.04872.x
26. Mithoowani S, Cervi A, Shah N, et al. Management of major bleeds in patients with immune thrombocytopenia. *J Thromb Haemost*. 2020;18(7):1783–1790. doi:10.1111/jth.14809
27. Fanouriakis A, Kostopoulou M, Alunno A, et al. 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. *Ann Rheum Dis*. 2019;78(6):736–745. doi:10.1136/annrheumdis-2019-215089
28. Zhu FX, Huang JY, Ye Z, Wen QQ, Wei JCC. Risk of systemic lupus erythematosus in patients with idiopathic thrombocytopenic purpura: A population-based cohort study. *Ann Rheum Dis*. 2020;79(6):793–799. doi:10.1136/annrheumdis-2020-217013
29. Jiang Y, Cheng Y, Ma S, et al. Systemic lupus erythematosus-complicating immune thrombocytopenia: From pathogenesis to treatment. *J Autoimmun*. 2022;132:102887. doi:10.1016/j.jaut.2022.102887
30. Neunert C, Terrell DR, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. *Blood Adv*. 2019;3(23):3829–3866. doi:10.1182/bloodadvances.2019000966
31. Sun L, Wang J, Shao L, et al. Dexamethasone plus oseltamivir versus dexamethasone in treatment-naïve primary immune thrombocytopenia: A multicentre, randomised, open-label, phase 2 trial. *Lancet Haematol*. 2021;8(4):e289–e298. doi:10.1016/S2352-3026(21)00030-2
32. Yang F, Lin J, Chen W. Post-translational modifications in T cells in systemic erythematosus lupus. *Rheumatology (Oxford)*. 2021;60(6):2502–2516. doi:10.1093/rheumatology/keab095
33. Mahévas M, Azzaoui I, Crickx E, et al. Efficacy, safety and immunological profile of combining rituximab with belimumab for adults with persistent or chronic immune thrombocytopenia: Results from a prospective phase 2b trial. *Haematologica*. 2020;106(9):2449–2457. doi:10.3324/haematol.2020.259481
34. Lucchini E, Zaja F, Bussell J. Rituximab in the treatment of immune thrombocytopenia: What is the role of this agent in 2019? *Haematologica*. 2019;104(6):1124–1135. doi:10.3324/haematol.2019.218883
35. Al-Ghazawi Z, Al-Farajat A, Toubasi AA, et al. Pancytopenia with aplastic anemia in systemic lupus erythematosus: Case series and literature review. *Rheumatol Int*. 2024;44(5):943–953. doi:10.1007/s00296-024-05585-6
36. Lally J, Boasman K, Brown L, et al. GATA-1: A potential novel biomarker for the differentiation of essential thrombocythemia and myelofibrosis. *J Thromb Haemost*. 2019;17(6):896–900. doi:10.1111/jth.14433
37. Santana IU, Dias B, Nunes EAS, Rocha FADC, Silva FS, Santiago MB. Visceral leishmaniasis mimicking systemic lupus erythematosus: Case series and a systematic literature review. *Semin Arthritis Rheum*. 2015;44(6):658–665. doi:10.1016/j.semarthrit.2014.12.004
38. Serin I, Bayir A, Goze H, Yokus O. A case report: Paroxysmal nocturnal hemoglobinuria and systemic lupus erythematosus association. *Int J Hematol Oncol*. 2021;10(1):IJH30. doi:10.2217/ijh-2020-0013
39. Frikha O, Ben Salah R, Mefteh A, et al. Agranulocytosis in systemic lupus erythematosus: A case report. *Clin Case Rep*. 2022;10(11):e6511. doi:10.1002/ccr3.6511
40. Sahbani H, Bahri M, Azza E, Ben Abdennebi Y, Aissaoui L. Atypical presentation of systemic lupus erythematosus in a 16-year-old boy: Case report. *Tunis Med*. 2024;102(10):725–729. doi:10.62438/tunismed.v102i10.5065
41. Horino T, Inotani S, Komori M, Ohnishi H, Terada Y. Neuropsychiatric systemic lupus erythematosus in a patient with pancytopenia and chronic schizophrenia requiring hospitalisation. *Intern Med*. 2025;64(1):147–151. doi:10.2169/internalmedicine.3202-23
42. Haidar G, Khalayli N, Drie T, Safiah MH, Kudsi M. Pancytopenia as a first presentation of late-onset systemic lupus erythematosus: A case report. *Ann Med Surg (Lond)*. 2024;86(5):3025–3028. doi:10.1097/MS9.0000000000001891
43. Brar Y, Yang NR, Poliyedath A. Hydralazine-induced lupus presenting with pancytopenia: A case report. *Cureus*. 2022;14(9):e29793. doi:10.7759/cureus.29793
44. Ungprasert P, Chowdhary VR, Davis MD, Makol A. Autoimmune myelofibrosis with pancytopenia as a presenting manifestation of systemic lupus erythematosus responsive to mycophenolate mofetil. *Lupus*. 2016;25(4):427–430. doi:10.1177/0961203315615221
45. Guevara NA, Flores Chang MM, Castelar J, Sequeira H, Berger J. A challenging diagnosis of febrile pancytopenia in a patient with a history of autoimmune disease. *Cureus*. 2023;15(3):e35956. doi:10.7759/cureus.35956
46. Abu Jheasha A, Alsharif T, Alwahsh R, Abumunshar A, Al-Ardah R, Abuturki A. Protein-losing enteropathy as the first presentation of systemic lupus erythematosus the first case reported in Palestine with systemic review. *Ann Med Surg (Lond)*. 2024;86(12):7458–7464. doi:10.1097/MS9.0000000000002733
47. Ameer MA, Tariq MA, Zain S, Kabir A, Khawaja M. A rare cause of pancytopenia in systemic lupus erythematosus (SLE) in a young patient. *Cureus*. 2024;16(6):e63032. doi:10.7759/cureus.63032
48. Wise LM, Stohl W. Belimumab and rituximab in systemic lupus erythematosus: A tale of two B cell-targeting agents. *Front Med (Lausanne)*. 2020;7:303. doi:10.3389/fmed.2020.00303
49. Serris A, Amoura Z, Canoui-Poitrine F, et al. Efficacy and safety of rituximab for systemic lupus erythematosus-associated immune cytopenias: A multicenter retrospective cohort study of 71 adults. *Am J Hematol*. 2018;93(3):424–429. doi:10.1002/ajh.24999
50. Khellaf M, Charles-Nelson A, Fain O, et al. Safety and efficacy of rituximab in adult immune thrombocytopenia: Results from a prospective registry including 248 patients. *Blood*. 2014;124(22):3228–3236. doi:10.1182/blood-2014-06-582346
51. Michel M, Terriou L, Roudot-Thoraval F, et al. A randomized and double-blind controlled trial evaluating the safety and efficacy of rituximab for warm auto-immune hemolytic anemia in adults (the RAIHA study). *Am J Hematol*. 2017;92(1):23–27. doi:10.1002/ajh.24570
52. Labrosse R, Barmettler S, Derfalvi B, et al. Rituximab-induced hypogammaglobulinemia and infection risk in pediatric patients. *J Allergy Clin Immunol*. 2021;148(2):523–532.e8. doi:10.1016/j.jaci.2021.03.041
53. Mo S, Li Y, He J, Lin L. Progress of rituximab in the treatment of systemic lupus erythematosus and lupus nephritis. *Front Med (Lausanne)*. 2024;11:1472019. doi:10.3389/fmed.2024.1472019
54. Boross P, Leusen JHW. Mechanisms of action of CD20 antibodies. *Am J Cancer Res*. 2012;2(6):676–690. PMID:23226614. PMCID:PMC3512181.
55. Junga Z, Stitt R, Tracy C, Keith M. Novel use of rituximab in macrophage activation syndrome secondary to systemic lupus erythematosus. *BMJ Case Rep*. 2017;2017:bcr-2017-221347. doi:10.1136/bcr-2017-221347
56. Serris A, Amoura Z, Terrier B, et al. Can rituximab be useful for treating SLE-associated immune cytopenias? Results from a retrospective multicentre study on 62 patients. *Blood*. 2016;128(22):870. doi:10.1182/blood.V128.22.870.870

57. Sans-Pola C, Danés I, Bosch JA, Marrero-Álvarez P, Cortés J, Agustí A. Off-label use of rituximab in patients with systemic lupus erythematosus with extrarenal disease activity: A retrospective study and literature review. *Front Med (Lausanne)*. 2023;10:1159794. doi:10.3389/fmed.2023.1159794
58. Rodrigues MM, Oliveira P, Simão C. 1009 rituximab in systemic lupus erythematosus in pediatric patients. *Pediatr Res*. 2010;68:502. doi:10.1203/00006450-201011001-01009
59. Duxbury B, Combescure C, Chizzolini C. Rituximab in systemic lupus erythematosus: An updated systematic review and meta-analysis. *Lupus*. 2013;22(14):1489–1503. doi:10.1177/0961203313509295
60. Zhang S, Jiang N, Wang L, et al. A modified regimen of low-dose rituximab therapy for patients with refractory immune thrombocytopenia associated with systemic lupus erythematosus. *Ther Adv Chronic Dis*. 2021;12:20406223211048643. doi:10.1177/20406223211048643
61. Harris EM, Hillier K, Al-Samkari H, Berbert L, Grace RF. Response to rituximab in children and adults with immune thrombocytopenia (ITP). *Res Pract Thromb Haemost*. 2021;5(6):e12587. doi:10.1002/rth2.12587
62. Tieu J, Smith RM, Gopaluni S, et al. Rituximab-associated hypogammaglobulinemia in autoimmune disease. *Front Immunol*. 2021;12:671503. doi:10.3389/fimmu.2021.671503
63. Zhang X, Rao M, Xu G. An update on the incidence, risk factors and mechanisms of rituximab-associated neutropenia. *Arch Med Sci*. 2022;20(2):494–505. doi:10.5114/aoms/152174
64. Gökten DB, Kavak FY, Mercan R. Could rituximab be a silver lining in refractory bone marrow fibrosis caused by lupus? *Arch Rheumatol*. 2024;39(3):476–478. doi:10.46497/ArchRheumatol.2024.10564
65. Li A, Guo F, Pan Q, et al. Mesenchymal stem cell therapy: Hope for patients with systemic lupus erythematosus. *Front Immunol*. 2021;12:728190. doi:10.3389/fimmu.2021.728190
66. Sui W, Hou X, Che W, et al. Hematopoietic and mesenchymal stem cell transplantation for severe and refractory systemic lupus erythematosus. *Clin Immunol*. 2013;148(2):186–197. doi:10.1016/j.clim.2013.05.014
67. Kirou KA, Dall'Era M, Aranow C, Anders HJ. Belimumab or anifrolumab for systemic lupus erythematosus? A risk–benefit assessment. *Front Immunol*. 2022;13:980079. doi:10.3389/fimmu.2022.980079
68. Singh JA, Shah NP, Mudano AS. Belimumab for systemic lupus erythematosus. *Cochrane Database Syst Rev*. 2021;2021(2):CD010668. doi:10.1002/14651858.CD010668.pub2
69. Levy RA, Gonzalez-Rivera T, Khamashta M, et al. 10 years of belimumab experience: What have we learnt? *Lupus*. 2021;30(11):1705–1721. doi:10.1177/09612033211028653
70. Fanouriakis A, Kostopoulou M, Andersen J, et al. EULAR recommendations for the management of systemic lupus erythematosus: 2023 update. *Ann Rheum Dis*. 2024;83(1):15–29. doi:10.1136/ard-2023-224762
71. Kalunian KC, Furie R, Morand EF, et al. A randomized placebo-controlled phase III extension trial of the long-term safety and tolerability of anifrolumab in active systemic lupus erythematosus. *Arthritis Rheumatol*. 2023;75(2):253–265. doi:10.1002/art.42392