

Arthroscopic vs open surgery for shoulder dislocation and instability: A network meta-analysis of treatment outcomes

Chenghong Wen^{A,D–F}, Qiang Hua^{B,C,F}, Wenduo Qian^{B,C,F}, Jide Su^{B,C,F}, Mingming Lei^{A,D–F}

Department of Sports Injury, Affiliated Sport Hospital of Chengdu Sport University, China

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation; D – writing the article; E – critical revision of the article; F – final approval of the article

Advances in Clinical and Experimental Medicine, ISSN 1899–5276 (print), ISSN 2451–2680 (online)

Adv Clin Exp Med. 2026;35(5):765–778

Address for correspondence

Chenghong Wen
E-mail: wenchenghong610@hotmail.com

Funding sources

None declared

Conflict of interest

None declared

Received on July 5, 2024

Reviewed on December 1, 2024

Accepted on July 24, 2025

Published online on April 7, 2026

Cite as

Wen C, Hua Q, Qian W, Su J, Lei M. Arthroscopic vs open surgery for shoulder dislocation and instability: A network meta-analysis of treatment outcomes. *Adv Clin Exp Med.* 2026;35(5):765–778. doi:10.17219/acem/208614

DOI

10.17219/acem/208614

Copyright

Copyright by Author(s)

This is an article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) (<https://creativecommons.org/licenses/by/3.0/>)

Abstract

Background. No comprehensive comparative research has been conducted to evaluate open Bankart (OB), arthroscopic Bankart (AB), open Latarjet (OL), and arthroscopic Latarjet (AL) simultaneously across all relevant clinical outcomes and parameters.

Objectives. To compare the efficacy of OB, AB, OL, and AL procedures in the treatment of shoulder dislocation.

Materials and methods. The databases PubMed, Embase, the Cochrane Library, and Web of Science were utilized for the literature search. The study evaluated recurrent instability, re-dislocation, apprehension, functional outcomes, and postoperative pain. The results were visually represented through network diagrams, forest plots, league tables, and rank probability plots to provide a comprehensive understanding of each outcome.

Results. Overall, 37 studies were included in the analysis. Individuals who underwent OL experienced a notably reduced risk of recurrent instability compared with those who underwent AB (random-effects model pooled relative risk (RR) = 0.34, 95% credible interval (95% CrI): 0.24–0.48) and OB (random-effects model pooled RR = 0.51, 95% CrI: 0.31–0.85). The risk of re-dislocation was also significantly lower for patients treated with OL compared with AB (pooled RR = 0.15, 95% CrI: 0.04–0.45). While not statistically significant, the OL procedure tended to have the lowest risk of apprehension and the highest Subjective Shoulder Value (SSV) score. Regarding postoperative pain, patients who underwent OB had the highest likelihood of attaining the lowest scores on the visual analogue scale (VAS). In addition, OL was associated with the highest probability of complications.

Conclusions. The open Latarjet procedure appears to offer superior shoulder stability; however, while functional outcomes for patients undergoing OL are likely to be non-inferior, the procedure is not significantly associated with reduced postoperative pain as measured with the VAS score. Additionally, the OL procedure is associated with an increased likelihood of complications. Consequently, it is essential to implement preventive measures to manage postoperative pain and address potential complications following OL procedure.

Key words: network meta-analysis, shoulder dislocation, Bankart, Open Latarjet, arthroscopic treatment

Highlights

- Open Latarjet procedure provides superior shoulder stability with significantly lower rates of recurrent instability and re-dislocation compared to Bankart techniques.
- Arthroscopic and open Bankart repairs show higher recurrence risk, while open Latarjet demonstrates the most favorable stability outcomes in shoulder dislocation treatment.
- Postoperative pain outcomes vary, with open Bankart associated with lower pain scores, while open Latarjet does not significantly reduce pain despite better stability.
- Open Latarjet carries a higher risk of complications, highlighting the need for careful patient selection and post-operative management strategies.

Introduction

The shoulder joint is the most frequently dislocated joint in the human body, with an incidence of approx. 23.96 per 100,000 individuals annually.¹ Its high mobility and relatively shallow glenoid cavity make it particularly susceptible to instability, especially following traumatic events.¹ Injuries to the static (e.g., labrum, ligaments) and dynamic (e.g., muscles) stabilizers of the shoulder can lead to instability, with anterior shoulder instability being the most common form, typically resulting from traumatic events.^{2,3} Conservative management is frequently selected for patients who are not candidates for surgery, despite the higher risk of recurrence.⁴ Surgical stabilization represents an effective treatment option when conservative management fails, providing durable fixation and a more definitive resolution of instability.⁵

Traditional open Bankart (OB) repair was historically regarded as the standard surgical treatment for shoulder stabilization.^{6,7} This technique has been shown to improve joint stability and is associated with low recurrence rates.^{8–11} However, despite its effectiveness, the open approach has certain disadvantages, including restricted external rotation and an increased risk of secondary osteoarthritis.¹²

Arthroscopic Bankart (AB) repair, first described in 1993, has gained increasing acceptance over the past decades owing to advances in arthroscopic techniques and improved surgical expertise.^{13–15} Compared with open procedures, arthroscopic repair offers several advantages, including smaller incisions, shorter operative time, reduced postoperative pain, and potentially fewer complications.^{16–18} However, some studies have reported higher recurrence rates of shoulder instability following AB repair compared with open repair.^{19,20}

The open Latarjet (OL) procedure is widely used for the management of anterior shoulder instability and is recognized for its effectiveness and reliability; however, it is associated with potential complications such as non-union and infection.^{21,22} Arthroscopic Latarjet (AL) has subsequently been introduced as a minimally invasive alternative. Emerging evidence indicates that AL achieves clinical outcomes comparable to OL, while potentially

offering smaller incisions, fewer complications, faster graft healing, earlier rehabilitation, and the ability to address concomitant intra-articular pathologies.^{23,24}

Numerous studies have directly compared pairs of the 4 surgical techniques – OB, AB, OL, and AL – for the management of shoulder instability.^{25–27} Furthermore, several meta-analyses have evaluated the comparative effectiveness of OL compared with AL in recurrent anterior shoulder instability, as well as OB vs AB in the management of Bankart lesions.^{28,29}

Currently, multiple surgical interventions are available for managing shoulder dislocations; however, no single treatment has been proven superior. Additionally, there is a scarcity of studies evaluating OB, AB, OL, and AL across all relevant clinical outcomes and parameters.

Objectives

The current network meta-analysis aimed to compare the OB, AB, OL, and AL procedures for the treatment of shoulder dislocation.

Materials and methods

Search strategy

To ensure a thorough examination of the available literature, 2 independent investigators (Q.H. and W.Q.) conducted a comprehensive search across multiple databases, including PubMed, Embase, the Cochrane Library, and Web of Science. The search was completed on August 17, 2023, to gather the most recent and relevant studies for the network meta-analysis. English search terms included “Shoulder Dislocation” OR “Dislocation, Shoulder” OR “Dislocations, Shoulder” OR “Shoulder Dislocations” OR “Glenohumeral Dislocation” OR “Dislocation, Glenohumeral” OR “Dislocations, Glenohumeral” OR “Glenohumeral Dislocations” OR “Glenohumeral Subluxation” OR “Glenohumeral Subluxations” OR “Subluxation, Glenohumeral” OR “Subluxations, Glenohumeral” AND “Bristow” OR “Latarjet” OR “Bankart” OR “iliac bone graft”

OR “Repair” OR “Putti-Platt” OR “Arthroscopies” OR “arthroscopic surgery” OR “Surgery” OR “Conservative” OR “nonoperative” OR “nonsurgical” OR “Physiotherapy” OR “Immobilization”. Disagreements concerning eligibility were resolved by another investigator (J.S.). Primary study selection was based on the titles and abstracts of the retrieved studies, followed by full-text screening. The present study adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines, ensuring a transparent and methodologically rigorous approach to the systematic review and meta-analysis process.³⁰

Eligibility criteria

The inclusion criteria for the study were as follows: 1) studies involving patients diagnosed with shoulder instability or dislocation; 2) studies in which patients underwent treatment with one of the 4 specified surgical procedures: OB, AB, OL, or AL; 3) evaluation of outcomes related to shoulder stability, functional outcomes, or postoperative pain; 4) study designs limited to randomized controlled trials (RCTs) and cohort studies.

The exclusion criteria were defined to ensure the relevance and quality of the included literature and included: 1) animal studies; 2) case reports; 3) studies with sample sizes of fewer than 10 participants; 4) studies not directly related to the topic, such as those involving patients without shoulder dislocation or instability, or those focusing on non-surgical treatments or procedures other than Bankart and Latarjet repairs; 5) editorial materials, conference abstracts, protocols, letters, guidelines, expert consensus documents, reviews, and meta-analyses, as these formats typically do not contain the level of detail or original data required for a systematic review and meta-analysis; 6) studies not published in English.

Outcome measures

In this analysis, the primary outcome of interest was shoulder stability, with specific measures including the incidence of recurrent instability, re-dislocation, and apprehension. The secondary outcomes encompassed functional outcomes and postoperative pain. Functional outcomes were measured using several standardized scales, including the Subjective Shoulder Value (SSV) score, reflecting the patient’s assessment of shoulder function; the American Shoulder and Elbow Surgeons (ASES) score, a comprehensive evaluation of shoulder function commonly used after shoulder surgery; the Rowe score, another scale specifically designed to assess shoulder function, with an emphasis on activities of daily living and sports; and forward flexion, a measure of range of motion indicating the extent to which the patient can raise the arm forward. Postoperative pain was quantified using the visual analogue scale (VAS).

Data extraction

The data extraction process from the included studies in the systematic review and network meta-analysis was comprehensive and included the following details: authors, year of publication, country, study design, population, intervention, sample size, male/female distribution, age (in years), body mass index (BMI, in kg/m²), glenoid bone loss, number of Hill–Sachs lesions, follow-up time (in months), and outcomes.

Quality assessment

In the systematic review and network meta-analysis, the quality assessment of the included RCTs was conducted using the modified Jadad scale.³¹ This scale allocates a total of 7 points, with RCT quality categorized as follows: 1–3 points indicating low quality and 4–7 points indicating high quality. For cohort studies, quality was evaluated using the modified Newcastle–Ottawa scale (NOS).³² The NOS assigns a total of 9 points, with quality ratings defined as follows: 0–3 points indicating poor quality, 4–6 points indicating fair quality, and 7–9 points indicating good quality.

Statistical analyses

A network meta-analysis was conducted using Stata v. 15.1 (StataCorp LLC, College Station, USA) and the gemtc 1.0.1 package in R v. 4.1.3 (R Foundation for Statistical Computing, Vienna, Austria). A Bayesian framework and a Markov chain Monte Carlo (MCMC) model were employed for the analysis. To ensure model convergence, trace plots and diagnostic tools were utilized. The number of initial iterations in the Monte Carlo simulations was determined based on achieving a potential scale reduction factor (PSRF) close to 1, indicating satisfactory convergence. Additional iterations were performed if necessary.

The bandwidth value in the density plots served as a quantitative measure, with smaller values suggesting a closer fit to the preset distribution. Iterations were continued until the bandwidth stabilized near 0, ensuring robust model performance. The model consisted of 4 chains, with an initial iteration count of 20,000, followed by an additional 50,000 iterations, and a step length of 1. In the presence of network relationships, the model assessed consistency and potential discrepancies between direct and indirect treatment comparisons. When the difference between the Deviance Information Criteria (DIC) of the consistency and inconsistency models was less than 5, it indicated agreement between direct and indirect comparisons. Network plots were constructed for each outcome measure. Analyses were conducted using a random-effects model. For the Rowe score, SSV score, forward flexion, ASES score, and VAS, the weighted

mean difference (WMD) and 95% credible intervals (95% CrIs) for different surgical interventions were reported. For recurrent instability, apprehension, re-dislocation, and complications, the relative risk (RR) values and 95% CrIs for different surgical interventions were reported. All direct and indirect comparisons of WMD or RR values with their corresponding 95% CrIs were presented in forest plots. The ranking of surgical interventions was predicted using a ranking plot, in which the probability of each intervention being ranked in the *n*th position was presented as a bar chart, with the horizontal axis indicating the corresponding rank.

Continuous variables were analyzed using the change from baseline as the final analysis outcome to reduce statistical errors arising from initial population imbalances in some cohort studies. An adjusted funnel plot was employed to evaluate potential publication bias. This approach is particularly relevant in the context of network meta-analyses, where traditional funnel plots may not adequately capture the nuances of multiple intervention comparisons. By using the adjusted funnel plot, symmetry of effect sizes across studies can be assessed more accurately, thereby providing a more robust evaluation of potential bias.

Results

Characteristics of the included studies

From the 4 databases, 8,402 studies were retrieved. After excluding duplicates, 5,982 studies remained for screening based on titles and abstracts, followed by full-text screening. Ultimately, 37 studies were included in this network meta-analysis, comprising 5 RCTs and 32 cohort studies.^{25–27,33–66} Figure 1 illustrates the flow chart of the eligible study screening process. Of the included studies, 32 were of high quality and 5 were of fair quality. These studies were published between 1997 and 2023. The characteristics of the included studies are presented in Supplementary Table 1.

Shoulder stability

Recurrent instability

Information on recurrent instability was provided in 21 studies involving 2,300 patients, in which OB, AB, OL, and AL were compared. Arthroscopic Bankart was directly compared with OB, AL, and OL. There were more studies for the direct comparison between AB and OL (Fig. 2A). The forest plot analysis using a random-effects




Fig. 1. Flowchart of eligible study screening

Table 1. League table of different procedures for various outcomes analyzed by the random-effect model

Outcomes/procedures		AB	AL	OB	OL
Recurrent instability	AB	AB	0.72 (0.22, 2.52)	0.67 (0.42, 1.04)	0.34 (0.24, 0.48)
	AL	1.38 (0.38, 4.57)	AL	0.94 (0.24, 3.3)	0.47 (0.13, 1.57)
	OB	1.47 (0.95, 2.36)	1.07 (0.3, 4.18)	OB	0.51 (0.31, 0.85)
	OL	2.95 (2.07, 4.18)	2.13 (0.64, 7.77)	1.99 (1.17, 3.34)	OL
Apprehension	AB	AB	0.76 (0.13, 3.4)	1.88 (0.24, 19.33)	0.57 (0.18, 1.45)
	AL	1.31 (0.29, 7.75)	AL	2.49 (0.21, 50.16)	0.75 (0.22, 2.68)
	OB	0.53 (0.05, 4.16)	0.4 (0.02, 4.83)	OB	0.3 (0.02, 2.79)
	OL	1.75 (0.69, 5.64)	1.33 (0.37, 4.56)	3.33 (0.36, 48.3)	OL
Re-dislocation	AB	AB	0.25 (0, 28.83)	0.65 (0.18, 2.09)	0.15 (0.04, 0.45)
	AL	3.96 (0.03, 524.99)	AL	2.55 (0.02, 360.41)	0.59 (0.01, 62.89)
	OB	1.55 (0.48, 5.55)	0.39 (0, 51.8)	OB	0.23 (0.04, 1.04)
	OL	6.71 (2.25, 27.26)	1.73 (0.02, 183.9)	4.34 (0.96, 24.12)	OL
Complications	AB	AB	0.98 (0.27, 4.77)	1.27 (0.39, 4.19)	2.08 (0.97, 6.26)
	AL	1.02 (0.21, 3.71)	AL	1.31 (0.18, 6.78)	2.12 (0.77, 6.53)
	OB	0.79 (0.22, 2.67)	0.76 (0.15, 5.41)	OB	1.64 (0.48, 8.38)
	OL	0.48 (0.15, 1.01)	0.47 (0.14, 1.32)	0.61 (0.12, 2.13)	OL
Rowe scores	AB	AB	-2.36 (-12.83, 8.38)	5.92 (-5.56, 16.63)	3.77 (-2.49, 10.28)
	AL	2.36 (-8.58, 12.83)	AL	8.3 (-7.65, 23.12)	6.22 (-3.79, 15.77)
	OB	-5.92 (-16.63, 5.46)	-8.3 (-23.12, 7.65)	OB	-2.14 (-14.5, 11.14)
	OL	-3.77 (-10.28, 2.49)	-6.12 (-15.77, 3.79)	2.14 (-11.14, 14.5)	OL
SSV score [%]	AB	AB	3.83 (-32.47, 40.42)	6.97 (-13.12, 27.37)	-
	AL	-3.83 (-40.42, 32.47)	AL	3.1 (-27.24, 33.4)	-
	OL	-6.97 (-27.37, 13.12)	-3.1 (-33.4, 27.24)	OL	-
Forward flexion	AB	AB	-14.36 (-32.77, 3.36)	-1.68 (-12.05, 7.64)	-
	AL	14.36 (-3.36, 32.77)	AL	12.6 (-2.54, 27.61)	-
	OL	1.68 (-7.65, 12.05)	-12.6 (-27.61, 2.58)	OL	-
ASES score	AB	AB	-22.88 (-47.99, 2.49)	-13.8 (-31.3, 3.96)	-
	AL	22.88 (-2.49, 47.99)	AL	9.1 (-8.8, 27.07)	-
	OL	13.8 (-3.96, 31.3)	-9.1 (-27.07, 8.8)	OL	-
VAS	AB	AB	-0.40 (-3.24, 2.44)	0.29 (-1.74, 2.31)	-
	OB	0.4 (-2.44, 3.24)	OB	0.69 (-2.81, 4.19)	-
	OL	-0.29 (-2.31, 1.75)	-0.69 (-4.19, 2.81)	OL	-

SSV – subjective shoulder value; ASES – American Shoulder and Elbow Surgeons; VAS – visual analogue scale; OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet. Values in brackets are 95% credible intervals (95 CrIs).

model revealed that OL had a lower risk of recurrent instability than AB (pooled relative risk (RR) = 0.33, 95% CrI: 0.22–0.49) (Fig. 3). The league table demonstrated that patients treated with OL had a significantly lower risk of recurrent instability than those treated with AB (random-effects model pooled RR = 0.34, 95% CrI: 0.24–0.48) and OB (random-effects model pooled RR = 0.51, 95% CrI: 0.31–0.85) (Table 1). The rank probabilities showed that OL was most likely to be the optimal procedure with regard to recurrent instability (Table 2). In the subgroup analysis restricted to primary surgical procedures, the pooled results analyzed using a random-effects model demonstrated that patients undergoing OL exhibited a significantly reduced risk of recurrent

instability compared with those treated with arthroscopic AB (pooled RR = 0.39, 95% CrI: 0.20–0.60) (Supplementary Table 2).

Re-dislocation

Nineteen studies involving 2,318 patients assessed the risk of re-dislocation after AB, OB, AL, and OL. Arthroscopic Bankart was directly compared with OB and OL, and OL was directly compared with OB and AL. More studies compared AB and OL (Fig. 2B). Figure 4, based on the random-effects model analysis, also showed a lower risk of re-dislocation in patients treated with OL compared with AB (pooled RR = 0.13, 95% CrI: 0.03–0.43). The league table




Fig. 2. Network plots of different procedures for shoulder stability. A. recurrent instability; B. re-dislocation; C. apprehension; D. complications
OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet.




Fig. 3. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (recurrent instability)
OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.

analysis using a random-effects model found that the risk of re-dislocation after OL was significantly lower than that after AB (pooled RR = 0.15, 95% CrI: 0.04–0.45) (Table 1).

The rank probabilities showed that OL had the highest likelihood of being the optimal operation with regard to re-dislocation (Table 2). Based on the subgroup analysis




Fig. 4. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (re-dislocation)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.

of primary surgeries, OL was associated with a lower likelihood of re-dislocation compared with AB (random-effects model pooled RR = 0.14, 95% CrI: 0.03–0.39) and OB (random-effects model pooled RR = 0.11, 95% CrI: 0.01–0.66) (Supplementary Table 2).

Apprehension

The risk of apprehension was evaluated in 10 studies including 1,195 patients. Arthroscopic Bankart was directly compared with OL and OB, and AL was directly compared with OL. More studies compared AB and OL (Fig. 2C). The forest plot analysis using a random-effects model demonstrated that no significant difference was observed in the risk of apprehension among these 4 operations (Fig. 5). Similarly, no significant difference was observed in the risk of apprehension among these 4 operations in the league table analysis using a random-effects model (Table 1). The rank probabilities showed that patients undergoing OL were most likely to have the lowest risk of apprehension (Table 2).

Functional outcomes

SSV score

Three studies involving 500 patients investigated the SSV score, including AB, OL, and AL. No significant difference in the SSV score was found among these surgical treatments based on the forest plot and league table (Table 1). The rank probabilities suggested that OL was most likely to be associated with the highest SSV score (Table 2). Based on the subgroup analysis of primary surgeries, no significant difference in the SSV score was observed among these surgical treatments (Supplementary Table 2).

ASES score

Two studies involving 952 patients assessed the ASES score, including AB, OL, and AL. No significant difference was observed in the ASES score among these surgical treatments based on the league table analysis using

Table 2. Rank probabilities of different procedures for various outcomes analyzed using the random-effect model

Outcomes/procedures		[1]	[2]	[3]	[4]
Recurrent instability	AB	0.67388	0.313325	0.012795	0
	AL	0.295145	0.24836	0.35018	0.106315
	OB	0.03096	0.43655	0.52601	0.00648
	OL	0.000015	0.001765	0.111015	0.887205
Apprehension	AB	0.167745	0.54864	0.226875	0.05674
	AL	0.14133	0.24713	0.35775	0.25379
	OB	0.679765	0.132705	0.07371	0.11382
	OL	0.01116	0.071525	0.341665	0.57565
Re-dislocation	AB	0.579025	0.359475	0.06113	0.00037
	AL	0.25713	0.090375	0.25431	0.398185
	OB	0.163505	0.53854	0.280285	0.01767
	OL	0.00034	0.01161	0.404275	0.583775
Complications	AB	0.01209	0.18025	0.46991	0.33775
	AL	0.05388	0.24929	0.248675	0.448155
	OB	0.20487	0.328335	0.253815	0.21298
	OL	0.72916	0.242125	0.0276	0.001115
Rowe scores	AB	0.01475	0.15198	0.56608	0.26719
	AL	0.046865	0.10945	0.201175	0.64251
	OB	0.62036	0.20063	0.10547	0.07354
	OL	0.318025	0.53794	0.127275	0.01676
SSV score [%]	AB	0.160565	0.313975	0.52546	–
	AL	0.38385	0.23618	0.37997	
	OL	0.455585	0.449845	0.09457	
Forward flexion	AB	0.70733	0.25892	0.03375	--
	AL	0.025975	0.035085	0.93894	
	OL	0.266695	0.705995	0.02731	
ASES score	AB	0.934345	0.04722	0.018435	–
	AL	0.02623	0.0955	0.87827	
	OL	0.039425	0.85728	0.103295	
VAS	AB	0.235695	0.52824	0.236065	–
	OB	0.263845	0.18881	0.547345	
	OL	0.50046	0.28295	0.21659	

SSV – subjective shoulder value; ASES – American Shoulder and Elbow Surgeons; VAS – visual analogue scale; OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet.

[1] – the probability that the procedure ranks 1st (i.e., performs the best) for that outcome; [2] – the probability that the procedure ranks 2nd for that outcome; [3] – the probability that the procedure ranks 3rd for that outcome; [4] – the probability that the procedure ranks 4th (i.e., performs the worst) for that outcome.

a random-effects model (Table 1). The rank probabilities suggested that AB was most likely to be the best surgery with regard to the ASES score (Table 2).

Rowe score

The Rowe score was evaluated in 11 studies involving 1,152 patients, and comparisons were made among OB, AB, OL, and AL. No significant difference in the Rowe score was found among these operations when the analysis was conducted using a random-effects model (Fig. 6). Similarly, no significant difference in the Rowe score

was observed among these operations in the league table analysis using a random-effects model (Table 1). The rank probabilities showed that OB had the highest likelihood of being the optimal procedure with regard to the Rowe score (Table 2).

Forward flexion

Three studies involving 311 patients provided data on forward flexion, including AB, OL, and AL. No significant difference in forward flexion was observed among the 3 operations in the forest plot analysis using




Fig. 5. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (apprehension)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.




Fig. 6. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (Rowe score)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.

a random-effects model (Fig. 7). Similarly, no significant difference in forward flexion was observed among the 3 operations in the league table analysis using a random-effects model (Table 1). The rank probabilities showed that AL was most likely to be associated with the greatest forward flexion (Table 2).

Postoperative pain

Information on the VAS score was provided in 3 studies involving 1,294 patients, including AB, OL, and OB. When the analysis was conducted using a random-effects model, no significant difference was observed in the VAS score




Fig. 7. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (forward flexion)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.




Fig. 8. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (visual analogue scale (VAS) score)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.

among AB, OL, and OB (Fig. 8). Similarly, no significant difference in the VAS score was observed among AB, OL, and OB in the league table analysis using a random-effects model (Table 1). The rank probabilities demonstrated that OB had the highest probability of achieving the lowest VAS score (Table 2).

Complications

A total of 18 studies comprising 1,902 patients were included in the analysis. Complications were assessed across 4 surgical approaches: AB, OB, OL, and AL. Direct comparisons were made between OL and each of the other 3 surgical methods, with specific direct comparisons existing between OL and AB. The network plot indicated a larger node and a thicker connecting line between AB and OL, suggesting that more studies directly compared these 2 interventions and that the sample size for these comparisons was relatively larger (Fig. 2D).

No significant difference in complications was observed among the 4 methods in the forest plot analysis using a random-effects model (Fig. 9). Similarly, no significant difference in complications was observed among the 4 techniques in the league table analysis using a random-effects model (Table 1). The rank probabilities demonstrated that OL had the highest probability of complications (Table 2). The funnel plots for publication bias assessment are shown in Supplementary Fig. 1.

Discussion

In this network meta-analysis, a comparison of the pooled data regarding the clinical efficacy of the OB, AB, OL, and AL procedures for shoulder dislocation and instability was conducted. Open Latarjet was most likely to have the lowest risk of recurrent instability and re-dislocation. Although not statistically significant, OL was most likely to have the lowest risk of apprehension and the highest SSV score. However, OL had the highest probability of complications. In addition, with regard to postoperative pain, the OL procedure was not necessarily associated with a significantly lower VAS score, indicating that it might not be the best option in terms of postoperative pain management.

The results of the network meta-analysis indicate that the OL procedure may offer superior outcomes in terms of shoulder stability and function for patients with shoulder dislocation and instability. However, the analysis also highlights a critical area for improvement: the management of postoperative pain following the OL procedure. In a meta-analysis conducted by Wang et al.,²⁹ a comparative analysis of OB compared with AB repairs for Bankart lesions revealed that while the OB method provided superior shoulder stability, it was accompanied by limitations in shoulder mobility.

Conversely, the AB technique, although less invasive, was associated with better preservation of motion. In another




Fig. 9. Forest plots of different procedures for shoulder stability analyzed with the random-effect model analysis (complications)

OB – open Bankart; AB – arthroscopic Bankart; OL – open Latarjet; AL – arthroscopic Latarjet; CrI – credibility interval.

meta-analysis, a comprehensive evaluation was conducted to compare the clinical efficacy, as well as the rates of postoperative revisions and complications, between OL and AL. The study found no significant differences in the majority of the assessed outcomes between the 2 procedures. However, the AL approach was associated with a lower score on the Western Ontario Shoulder Instability Index, indicating better shoulder function or less instability. Additionally, the AL procedure exhibited a higher revision rate, suggesting a greater likelihood of patients requiring additional surgeries.²⁸

Nonetheless, the efficacy of OB, AB, OL, and AL has not been compared in patients with shoulder dislocation and instability in existing studies. To facilitate understanding and decision-making among the 4 methods when considering therapeutic options for shoulder dislocation, the present network meta-analysis focused on shoulder stability, functional outcomes, and postoperative pain following these procedures.

In terms of shoulder stability, the OL procedure was indicated to have the highest likelihood of being the most effective repair option. In a meta-analysis encompassing 795 shoulders, the OL procedure demonstrated a significantly lower risk of recurrence compared with AB repair.⁶⁷

As reported by Rollick et al.,⁶⁸ the estimated re-dislocation rate was notably higher at 15.1% for patients who had undergone AB, whereas it was considerably lower at 2.7% for those who had OL procedure. In terms of functional outcomes, OL may also demonstrate a relative advantage.

According to the findings by Bliven et al.,⁶⁹ a higher percentage of patients who underwent the Latarjet procedure were able to successfully return to work, sports, and throwing activities compared with those who had Bankart repair. Consistently, a study has shown that the Latarjet procedure for anterior shoulder instability leads to excellent long-term functional outcomes and a high rate of return to sports.⁷⁰ However, patients might experience unsatisfactory postoperative pain outcomes and complications following the OL procedure, which could be attributed to the larger incision associated with the surgery. Arthroscopic surgery generally offers the benefits of a lower nonunion rate and the potential for quicker recovery due to its minimally invasive nature. The drawbacks of the AL procedure are also evident. It requires longer operative time and a longer learning curve for surgeons.⁷¹

Analysis of the learning curve indicated that the initial cohort experienced longer operative times and higher complication rates compared with the subsequent group,

potentially accounting for the increased incidence of revision surgeries required for the AL procedure.⁷² Only surgeons expected to perform the AL procedure at high volume should consider adopting this technique.^{73,74} Additionally, it has been shown that the direct costs associated with the AL procedure are significantly higher, amounting to approximately double the cost of the OL procedure, with figures of 2,335 EUR for AL compared with 1,040 EUR for OL.⁷⁵

Several factors may contribute to the effectiveness of the OL procedure. The OL procedure provides a dual stabilizing effect by increasing the anterior glenoid width and creating a sling effect with the conjoint tendon, which helps stabilize the joint.⁷⁶ Correct graft positioning is critical to the success of the OL procedure. It has been suggested that OL might offer better control over graft placement, which could lead to improved long-term outcomes.⁷⁷

Limitations of the study

Several limitations of this study should be acknowledged. First, there was a lack of literature on some outcomes, which may have affected the accuracy of the results. Second, this study focused on the general population, and further research was not conducted in specific populations such as athletes and males or females.⁵¹ In addition, some of the included studies were cohort studies; therefore, change scores were used for continuous data, and studies that provided only endpoint data were not included in this analysis. Third, including only English-language literature may result in language bias. Fourth, there are differences between primary operations and re-operations, and these differences could potentially influence the outcomes of interest. However, due to limitations in the included literature, such as the lack of detailed data on re-operations, we were unable to conduct a separate analysis for re-operations. This precludes definitive conclusions regarding the comparative effectiveness of the 4 surgical approaches for shoulder dislocation and instability in the context of re-operations. Fifth, we acknowledge that glenoid bone loss, a significant factor affecting shoulder joint stability and surgical outcomes, was not controlled for in our screening process. Although glenoid bone loss was included as a variable in Supplementary Table 1, the substantial amount of missing data across the reviewed studies prevented us from accounting for it in our analysis. This limitation may restrict the accuracy of our comparisons among different surgical approaches.

Conclusions

Open Latarjet was most likely to be the optimal procedure for shoulder stability, and patients undergoing OL may have non-inferior functional outcomes. Although the OL procedure may not be associated with significantly

lower VAS pain scores, it was associated with a higher rate of complications. Therefore, it is imperative to implement appropriate measures to mitigate postoperative pain and manage complications following OL procedure.

Supplementary data

The supplementary materials are available at <https://doi.org/10.5281/zenodo.15872256>. The package contains the following files:

Supplementary Table 1. Characteristics of the included studies.

Supplementary Table 2. League table of different procedures for various outcomes in patients with primary operation analyzed by the random-effect model.

Supplementary Fig. 1. A. Publication bias assessed by the funnel plot for recurrent instability; B. Publication bias assessed by the funnel plot for re-dislocation; C. Publication bias assessed by the funnel plot for apprehension; D. Publication bias assessed by the funnel plot for Rowe; E. Publication bias assessed by the funnel plot for complications.

Use of AI and AI-assisted technologies

Not applicable.

ORCID iDs

Chenghong Wen  <https://orcid.org/0009-0006-7136-3302>
 Wenduo Qian  <https://orcid.org/0009-0002-7371-4154>
 Mingming Lei  <https://orcid.org/0009-0006-4088-4236>
 Qiang Hua  <https://orcid.org/0000-0002-4582-0382>
 Jide Su  <https://orcid.org/0009-0007-5512-1837>

References

- Patrick CM, Snowden J, Eckhoff MD, et al. Epidemiology of shoulder dislocations presenting to United States emergency departments: An updated ten-year study. *World J Orthop.* 2023;14(9):690–697. doi:10.5312/wjo.v14.i9.690
- Stokes DJ, McCarthy TP, Frank RM. Physical therapy for the treatment of shoulder instability. *Phys Med Rehabil Clin North Am.* 2023;34(2):393–408. doi:10.1016/j.pmr.2022.12.006
- Christopher HW, Grainger AJ. Anterior shoulder instability. *Semin Musculoskelet Radiol.* 2022;26(5):546–557. doi:10.1055/s-0042-1756168
- Do W, Kim J, Lim J, Yoon T, Shin S, Chun Y. High failure rate after conservative treatment for recurrent shoulder dislocation without subjective apprehension on physical examination. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(1):178–184. doi:10.1007/s00167-022-07028-w
- Belk JW, Wharton BR, Houck DA, et al. Shoulder stabilization versus immobilization for first-time anterior shoulder dislocation: A systematic review and meta-analysis of level 1 randomized controlled trials. *Am J Sports Med.* 2023;51(6):1634–1643. doi:10.1177/03635465211065403
- Fares MY, Boufadel P, Daher M, Koa J, Khanna A, Abboud JA. Anterior shoulder instability and open procedures: History, indications, and clinical outcomes. *Clin Orthop Surg.* 2023;15(4):521. doi:10.4055/cios23018
- AlSomali K, Kholinne E, Van Nguyen T, et al. Outcomes and return to sport and work after open Bankart repair for recurrent shoulder instability: A systematic review. *Orthop J Sports Med.* 2021;9(10):23259671211026907. doi:10.1177/23259671211026907
- Monk AP, Crua E, Gatenby GC, et al. Clinical outcomes following open anterior shoulder stabilization for glenohumeral instability in the young collision athlete. *J Shoulder Elbow Surg.* 2022;31(7):1474–1478. doi:10.1016/j.jse.2021.12.013

9. Mancini MR, Arciero RA. Open Bankart repair. *Clin Sports Med.* 2024; 43(4):617–633. doi:10.1016/j.csm.2023.12.002
10. Bitar I, Bustos D, Marangoni L, Robles C, Gentile L, Bertiche P. Outcomes of open Bankart repair plus inferior capsular shift compared with Latarjet procedure in contact athletes with recurrent anterior shoulder instability. *Arch Bone Joint Surg.* 2023;11(1):39–46. doi:10.22038/abjs.2022.60208.2974
11. Neviasser AS, Benke MT, Neviasser RJ. Open Bankart repair for revision of failed prior stabilization: Outcome analysis at a mean of more than 10 years. *J Shoulder Elbow Surg.* 2015;24(6):897–901. doi:10.1016/j.jse.2014.11.036
12. Berendes T, Mathijssen N, Verburg H, Kraan G. The open-modified Bankart procedure: Long-term follow-up 'a 16–26-year follow-up study.' *Arch Orthop Trauma Surg.* 2018;138(5):597–603. doi:10.1007/s00402-017-2866-9
13. Hu B, Hong J, Zhu H, Yan S, Wu H. Arthroscopic Bankart repair versus conservative treatment for first-time traumatic anterior shoulder dislocation: A systematic review and meta-analysis. *Eur J Med Res.* 2023;28(1):260. doi:10.1186/s40001-023-01160-0
14. Rashid MS, Arner JW, Millett PJ, Sugaya H, Emery R. The Bankart repair: Past, present, and future. *J Shoulder Elbow Surg.* 2020;29(12):e491–e498. doi:10.1016/j.jse.2020.06.012
15. Vermeulen AE, Landman EBM, Veen EJD, Nienhuis S, Koorevaar CT. Long-term clinical outcome of arthroscopic Bankart repair with suture anchors. *J Shoulder Elbow Surg.* 2019;28(5):e137–e143. doi:10.1016/j.jse.2018.09.027
16. McQuivey KS, Brinkman JC, Tummala SV, Shaha JS, Tokish JM. Arthroscopic remplissage using knotless, all-suture anchors. *Arthrosc Techn.* 2022;11(4):e615–e621. doi:10.1016/j.eats.2021.12.015
17. Su BY, Yi SY, Peng T, Yi G, Zhang L. Comparison of arthroscopic surgery versus open surgical repair of the anterior talofibular ligament: A retrospective study of 80 patients from a single center. *Med Sci Monit.* 2020;27:e928526. doi:10.12659/msm.928526
18. Ahmed AF, Alzobi OZ, Hantouly AT, et al. Complications of elbow arthroscopic surgery: A systematic review and meta-analysis. *Orthop J Sports Med.* 2022;10(11):23259671221137863. doi:10.1177/23259671221137863
19. Hurley ET, Manjunath AK, Bloom DA, et al. Arthroscopic Bankart repair versus conservative management for first-time traumatic anterior shoulder instability: A systematic review and meta-analysis. *Arthroscopy.* 2020;36(9):2526–2532. doi:10.1016/j.arthro.2020.04.046
20. Haskel JD, Wang KH, Hurley ET, et al. Clinical outcomes of revision arthroscopic Bankart repair for anterior shoulder instability: A systematic review of studies. *J Shoulder Elbow Surg.* 2022;31(1):209–216. doi:10.1016/j.jse.2021.06.021
21. Werthel JD, Sabatier V, Schoch B, et al. Outcomes of the Latarjet procedure for the treatment of chronic anterior shoulder instability: Patients with prior arthroscopic Bankart repair versus primary cases. *Am J Sports Med.* 2020;48(1):27–32. doi:10.1177/0363546519888909
22. Longo UG, Loppini M, Rizzello G, Ciuffreda M, Maffulli N, Denaro V. Latarjet, Bristow, and Eden–Hybinette procedures for anterior shoulder dislocation: Systematic review and quantitative synthesis of the literature. *Arthroscopy.* 2014;30(9):1184–1211. doi:10.1016/j.arthro.2014.04.005
23. Cerciello S, Corona K, Morris BJ, Santagada DA, Maccauro G. Early outcomes and perioperative complications of the arthroscopic Latarjet procedure: Systematic review and meta-analysis. *Am J Sports Med.* 2019;47(9):2232–2241. doi:10.1177/0363546518783743
24. Griesser MJ, Harris JD, McCoy BW, et al. Complications and re-operations after Bristow–Latarjet shoulder stabilization: A systematic review. *J Shoulder Elbow Surg.* 2013;22(2):286–292. doi:10.1016/j.jse.2012.09.009
25. Maiotti M, De Vita A, De Benedetto M, et al. Clinical outcomes and recurrence rate of 4 procedures for recurrent anterior shoulder instability: ASA, remplissage, open, and arthroscopic Latarjet. A multicenter study. *J Shoulder Elbow Surg.* 2023;32(5):931–938. doi:10.1016/j.jse.2022.10.030
26. Saremi H, Saneii A, Goodarzi B. Midterm clinical results of Bankart repair, Bankart remplissage, and Latarjet procedures for treating recurrent anterior shoulder dislocation. *Adv Hum Biol.* 2021;11(Suppl 1): S22–S26. doi:10.4103/aihb.aih22_21
27. Woodmass JM, Wagner ER, Smith J, et al. Postoperative recovery comparisons of arthroscopic Bankart to open Latarjet for the treatment of anterior glenohumeral instability. *Eur J Orthop Surg Traumatol.* 2022;33(4):1357–1364. doi:10.1007/s00590-022-03265-4
28. Deng Z, Zheng Y, Su J, et al. Open versus arthroscopic Latarjet for recurrent anterior shoulder instability: A systematic review and meta-analysis. *Orthop J Sports Med.* 2023;11(5):23259671231174476. doi:10.1177/23259671231174476
29. Wang L, Liu Y, Su X, Liu S. A meta-analysis of arthroscopic versus open repair for treatment of Bankart lesions in the shoulder. *Med Sci Monit.* 2015;21:3028–3035. doi:10.12659/msm.894346
30. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. doi:10.1136/bmj.n71
31. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: Is blinding necessary? *Cont Clin Trials.* 1996;17(1):1–12. doi:10.1016/0197-2456(95)00134-4
32. Wells GA, Shea BJ, O'Connell D, Peterson J, Tugwell P. The Newcastle–Ottawa Scale (NOS) for Assessing the Quality of Non-Randomized Studies in Meta-Analysis. Ottawa, Canada: The Ottawa Hospital Research Institute; 2020. <https://ohri.ca/en/who-we-are/core-facilities-and-platforms/ottawa-methods-centre/newcastle-ottawa-scale>. Accessed March 6, 2026.
33. Paul RW, Reddy MP, Sonnier JH, et al. Increased rates of subjective shoulder instability after Bankart repair with remplissage compared to Latarjet surgery. *J Shoulder Elbow Surg.* 2023;32(5):939–946. doi:10.1016/j.jse.2022.11.001
34. Min KS, Wake J, Cruz C, et al. Surgical treatment of shoulder instability in active-duty service members with subcritical glenoid bone loss: Bankart vs Latarjet. *J Shoulder Elbow Surg.* 2023;32(4):771–775. doi:10.1016/j.jse.2022.10.011
35. Gul Y, Farooq MZ, I, Essa MA, Khan SU. Evaluation of arthroscopic Bankart repair and open Latarjet technique for treatment of recurrent shoulder dislocation. *Pak J Med Health Sci.* 2023;17(2):528–530. doi:10.53350/pjmhs2023172528
36. Genena A, Hashem M, Waly A, Hegazy MO. Open Latarjet versus arthroscopic Bankart repair for the treatment of traumatic anterior shoulder instability in high-demand patients with minimal glenoid bone loss. *Cureus.* 2023;15(4):e37127. doi:10.7759/cureus.37127
37. Waltenspül M, Ernstbrunner L, Ackermann J, Thiel K, Galvin JW, Wieser K. Long-term results and failure analysis of the open Latarjet procedure and arthroscopic Bankart repair in adolescents. *J Bone Joint Surg.* 2022;104(12):1046–1054. doi:10.2106/jbjs.21.01050
38. Kukkonen J, Elamo S, Flinkkilä T, et al. Arthroscopic Bankart versus open Latarjet as a primary operative treatment for traumatic antero-inferior instability in young males: A randomised controlled trial with 2-year follow-up. *Br J Sports Med.* 2022;56(6):327–333. doi:10.1136/bjsports-2021-104028
39. Girard M, Dalmas Y, Martinel V, Mansat P, Bonnevalie N. Arthroscopic Latarjet with cortical buttons versus open Latarjet with screws: A short-term comparative study. *Am J Sports Med.* 2022;50(12): 3326–3332. doi:10.1177/03635465221120076
40. Rai S, Tamang N, Sharma LK, et al. Comparative study of arthroscopic Bankart repair versus open Latarjet procedure for recurrent shoulder dislocation. *J Int Med Res.* 2021;49(4):3000605211007328. doi:10.1177/03000605211007328
41. Calvo E, Luengo G, Morcillo D, Foruria AM, Valencia M. Revision arthroscopic bankart repair versus arthroscopic Latarjet for failed primary arthroscopic stabilization with subcritical bone loss. *Orthop J Sports Med.* 2021;9(5):23259671211001809. doi:10.1177/23259671211001809
42. Maman E, Dolkart O, Krespi R, et al. A multicenter retrospective study with a minimum 5-year follow-up comparing arthroscopic Bankart repair and the Latarjet procedure. *Orthop J Sports Med.* 2020; 8(8):2325967120941366. doi:10.1177/2325967120941366
43. Elamo S, Selänne L, Lehtimäki K, et al. Bankart versus Latarjet operation as a revision procedure after a failed arthroscopic Bankart repair. *JSES Int.* 2020;4(2):292–296. doi:10.1016/j.jseint.2020.01.004
44. Ali J, Altintas B, Pulatkan A, Boykin RE, Aksoy DO, Bilsel K. Open versus arthroscopic Latarjet procedure for the treatment of chronic anterior glenohumeral instability with glenoid bone loss. *Arthroscopy.* 2020;36(4):940–949. doi:10.1016/j.arthro.2019.09.042

45. Xu Y, Wu K, Ma Q, et al. Comparison of clinical and patient-reported outcomes of three procedures for recurrent anterior shoulder instability: Arthroscopic Bankart repair, capsular shift, and open Latarjet. *J Orthop Surg Res*. 2019;14(1):326. doi:10.1186/s13018-019-1340-5
46. Liang Z, Li B, Muheremu A. Arthroscopic versus open bankart repair in patients with recurrent anterior dislocation of the shoulder: A ten year follow-up study. *Int J Clin Exp Med*. 2019;12(6):7814–7819. <https://e-century.us/files/ijcem/12/6/ijcem0089952.pdf>.
47. Jeon YS, Jeong HY, Lee DK, Rhee YG. Borderline glenoid bone defect in anterior shoulder instability: Latarjet procedure versus Bankart repair. *Am J Sports Med*. 2018;46(9):2170–2176. doi:10.1177/0363546518776978
48. Bah A, Lateur GM, Kouevidjin BT, et al. Chronic anterior shoulder instability with significant Hill–Sachs lesion: Arthroscopic Bankart with remplissage versus open Latarjet procedure. *Orthop Traumatol Surg Res*. 2018;104(1):17–22. doi:10.1016/j.otsr.2017.11.009
49. Zhu Y, Jiang C, Song G. Arthroscopic versus open Latarjet in the treatment of recurrent anterior shoulder dislocation with marked glenoid bone loss: A prospective comparative study. *Am J Sports Med*. 2017;45(7):1645–1653. doi:10.1177/0363546517693845
50. Kordasiewicz B, Malachowski K, Kicinski M, Chaberek S, Pomianowski S. Comparative study of open and arthroscopic coracoid transfer for shoulder anterior instability (Latarjet): Clinical results at short term follow-up. *Int Orthop*. 2017;41(5):1023–1033. doi:10.1007/s00264-016-3372-3
51. Zimmermann SM, Scheyerer MJ, Farshad M, Catanzaro S, Rahm S, Gerber C. Long-term restoration of anterior shoulder stability: A retrospective analysis of arthroscopic Bankart repair versus open Latarjet procedure. *J Bone Joint Surg Am*. 2016;98(23):1954–1961. doi:10.2106/JBJS.15.01398
52. Virk MS, Manzo RL, Cote M, et al. Comparison of time to recurrence of instability after open and arthroscopic Bankart repair techniques. *Orthop J Sports Med*. 2016;4(6):2325967116654114. doi:10.1177/2325967116654114
53. Cho NS, Yoo JH, Rhee YG. Management of an engaging Hill–Sachs lesion: Arthroscopic remplissage with Bankart repair versus Latarjet procedure. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(12):3793–3800. doi:10.1007/s00167-015-3666-9
54. Blonna D, Bellato E, Caranzano F, Assom M, Rossi R, Castoldi F. Arthroscopic Bankart repair versus open Bristow–Latarjet for shoulder instability: A matched-pair multicenter study focused on return to sport. *Am J Sports Med*. 2016;44(12):3198–3205. doi:10.1177/0363546516658037
55. Shymon SJ, Roocroft J, Edmonds EW. Traumatic anterior instability of the pediatric shoulder: A comparison of arthroscopic and open bankart repairs. *J Pediatr Orthop*. 2015;35(1):1–6. doi:10.1097/BPO.0000000000000215
56. Abouelsoud MM, Abdelrahman AA. Recurrent anterior shoulder dislocation with engaging Hill–Sachs defect: Remplissage or Latarjet? *Eur Orthop Traumatol*. 2015;6(3):151–156. doi:10.1007/s12570-015-0313-3
57. Bessière C, Trojani C, Carles M, Mehta SS, Boileau P. The open Latarjet procedure is more reliable in terms of shoulder stability than arthroscopic Bankart repair. *Clin Orthop Relat Res*. 2014;472(8):2345–2351. doi:10.1007/s11999-014-3550-9
58. Hovelius L, Vikerfors O, Olofsson A, Svensson O, Rahme H. Bristow–Latarjet and Bankart: A comparative study of shoulder stabilization in 185 shoulders during a seventeen-year follow-up. *J Shoulder Elbow Surg*. 2011;20(7):1095–1101. doi:10.1016/j.jse.2011.02.005
59. Mahiroğulları M, Ozkan H, Akyüz M, Uğraş AA, Güney A, Kuşkuçcu M. Comparison between the results of open and arthroscopic repair of isolated traumatic anterior instability of the shoulder. *Acta Orthop Traumatol Turc*. 2010;44(3):180–185. doi:10.3944/AOTT.2010.2289
60. Tjoumakaris FP, Abboud JA, Hasan SA, Ramsey ML, Williams GR. Arthroscopic and open Bankart repairs provide similar outcomes. *Clin Orthop Relat Res*. 2006;446:227–232. doi:10.1097/01.blo.0000205883.73705.19
61. Bottoni CR, Smith EL, Berkowitz MJ, Towle RB, Moore JH. Arthroscopic versus open shoulder stabilization for recurrent anterior instability: A prospective randomized clinical trial. *Am J Sports Med*. 2006;34(11):1730–1737. doi:10.1177/0363546506288239
62. Wang C, Ghalambor N, Zarins B, Warner JJP. Arthroscopic versus open Bankart repair: Analysis of patient subjective outcome and cost. *Arthroscopy*. 2005;21(10):1219–1222. doi:10.1016/j.arthro.2005.07.004
63. Kim SH, Ha KI, Kim SH. Bankart repair in traumatic anterior shoulder instability. *Arthroscopy*. 2002;18(7):755–763. doi:10.1053/jars.2002.31701
64. Sperber A, Hamberg P, Karlsson J, Swärd L, Wredmark T. Comparison of an arthroscopic and an open procedure for posttraumatic instability of the shoulder: A prospective, randomized multicenter study. *J Shoulder Elbow Surg*. 2001;10(2):105–108. doi:10.1067/mse.2001.112019
65. Karlsson J, Magnusson L, Ejerhed L, Hultenheim I, Lundin O, Kartus J. Comparison of open and arthroscopic stabilization for recurrent shoulder dislocation in patients with a Bankart lesion. *Am J Sports Med*. 2001;29(5):538–542. doi:10.1177/03635465010290050201
66. Geiger DF, Hurley JA, Tovey JA, Rao JP. Results of arthroscopic versus open Bankart suture repair. *Clin Orthop Relat Res*. 1997;337:111–117. doi:10.1097/00003086-199704000-00013
67. An VVG, Sivakumar BS, Phan K, Trantalís J. A systematic review and meta-analysis of clinical and patient-reported outcomes following two procedures for recurrent traumatic anterior instability of the shoulder: Latarjet procedure vs Bankart repair. *J Shoulder Elbow Surg*. 2016; 25(5):853–863. doi:10.1016/j.jse.2015.11.001
68. Rollick N, Ono Y, Kurji HM, et al. Long-term outcomes of the Bankart and Latarjet repairs: A systematic review. *Open Access J Sports Med*. 2017;8:97–105. doi:10.2147/oajsm.s106983
69. Bliven KCH, Parr GP. Outcomes of the Latarjet procedure compared with Bankart repair for recurrent traumatic anterior shoulder instability. *J Athl Train*. 2018;53(2):181–183. doi:10.4085/1062-6050-232-16
70. Hurley ET, Jamal MS, Ali ZS, Montgomery C, Pauzenberger L, Mullett H. Long-term outcomes of the Latarjet procedure for anterior shoulder instability: A systematic review of studies at 10-year follow-up. *J Shoulder Elbow Surg*. 2019;28(2):e33–e39. doi:10.1016/j.jse.2018.08.028
71. Hurley ET, Lim Fat D, Farrington SK, Mullett H. Open versus arthroscopic Latarjet procedure for anterior shoulder instability: A systematic review and meta-analysis. *Am J Sports Med*. 2019;47(5):1248–1253. doi:10.1177/0363546518759540
72. Bøe B, Støen RØ, Blich I, Moatshe G, Ludvigsen TC. Learning curve for arthroscopic shoulder Latarjet procedure shows shorter operating time and fewer complications with experience. *Arthroscopy*. 2022;38(8):2391–2398. doi:10.1016/j.arthro.2022.01.042
73. Gendre P, Thélu CE, d'Ollonne T, Trojani C, Gonzalez JF, Boileau P. Coracoid bone block fixation with cortical buttons: An alternative to screw fixation? *Orthop Traumatol Surg Res*. 2016;102(8):983–987. doi:10.1016/j.otsr.2016.06.016
74. Valsamis EM, Kany J, Bonnevalle N, et al. The arthroscopic Latarjet: A multisurgeon learning curve analysis. *J Shoulder Elbow Surg*. 2020; 29(4):681–688. doi:10.1016/j.jse.2019.10.022
75. Randelli P, Fossati C, Stoppani C, Evola FR, De Girolamo L. Open Latarjet versus arthroscopic Latarjet: Clinical results and cost analysis. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(2):526–532. doi:10.1007/s00167-015-3978-9
76. Çağlar C, Akçaalan S, Akbulut B, Kengil MC, Uğurlu M, Doğan M. Open Latarjet reduces residual apprehension, re-dislocation and possibility of dislocation arthropathy compared to arthroscopic Bankart repair despite greater bipolar bone loss in anterior glenohumeral instability. *JSES Int*. 2024;8(6):1175–1181. doi:10.1016/j.jseint.2024.08.181
77. Vuletić F, Bøe B. Current trends and outcomes for open vs arthroscopic Latarjet. *Curr Rev Musculoskelet Med*. 2024;17(5):136–143. doi:10.1007/s12178-024-09889-9